

เซต \mathcal{A} ในปริภูมิสองโครงสร้างเล็กสุด

\mathcal{A} -Sets in Biminimal Structure Spaces

ชนิกา กุลค้อ¹, จีระนันท์ คำภักดี^{2*}, ดรุณี บุญชารี³

Chanika Kulkhor¹, Jeeranunt Khampakdee^{2*}, Daruni Boonchari³

Received: 20 December 2017; Accepted: 9 February 2018

บทคัดย่อ

งานวิจัยนี้จะนำเสนอแนวคิดเกี่ยวกับเซต \mathcal{A} ในปริภูมิสองโครงสร้างเล็กสุดและตรวจสอบคุณสมบัติบางประการของเซต \mathcal{A} รวมทั้งฟังก์ชันต่อเนื่อง \mathcal{A} ในปริภูมิสองโครงสร้างเล็กสุด

คำสำคัญ : เซต \mathcal{A} , ฟังก์ชันต่อเนื่อง \mathcal{A}

Abstract

In this article, we introduce the concepts of \mathcal{A} -sets in biminimal structure spaces and investigate some of their properties. Moreover, the notions of \mathcal{A} -sets and \mathcal{A} -continuous functions in biminimal structure spaces were studied.

Keywords : \mathcal{A} -set, \mathcal{A} -continuous function.

Introduction

In 1972, J. Dugundji⁷ introduced the concepts of regular closed sets in topological spaces. Let (X, τ) be a topological space and $A \subseteq X$, then A is called regular closed if and only if $A = Cl(Int(A))$. In 1986, J. Tong²¹ introduced the concepts and properties of \mathcal{A} -sets in topological spaces. Let A be a subset of a topological space (X, τ) then A is an \mathcal{A} -set in (X, τ) if $A = U \cap B$ when U is open and B is regular closed in (X, τ) . In addition, J. Tong²¹ introduced the concepts of \mathcal{A} -continuous functions from a topological space (X, τ) to a topological space (Y, \mathcal{U}) . Let f be a function from X to Y , then f is \mathcal{A} -continuous function if and only if the inverse image of each open set in Y is an \mathcal{A} -set in X . In 1990, M. Ganster, and Reilly, I. L.¹⁰ improved J. Tong's decomposition result and provided a decomposition of \mathcal{A} -continuous. In 2000, the concepts of minimal structure spaces were introduced by V. Popa and T. Noiri¹⁸. A pair (X, m_x) is a minimal structure space if and only if $X \neq \emptyset$ and m_x is family of

$P(X)$ with $\emptyset, X \in m_x$. Moreover, they also introduced the concepts of m_x -open sets and m_x -closed sets in minimal structure spaces. In 1963, J. C. Kelly⁹ introduced the concepts of bitopological spaces which consist of a nonempty set and two topological spaces. In 2010, C. Boonpok² introduced the concepts of biminimal structure spaces which consist of a nonempty set and two minimal structures. Furthermore, C. Boonpok² defined $m_x^1 m_x^2$ -closed sets in biminimal structure spaces and the complement of $m_x^1 m_x^2$ -closed sets is call $m_x^1 m_x^2$ -open sets. In 2010, C. Boonpok [4] defined (i, j) m_x -regular open sets in biminimal structure spaces and he also defined (i, j) m_x -regular closed sets as complement of (i, j) m_x -regular open sets for $i, j = 1, 2$ and $i \neq j$.

In this article we introduce the concepts of \mathcal{A} -sets in biminimal structure spaces and \mathcal{A} -continuous functions in biminimal structure spaces. Also, we study some properties of \mathcal{A} -sets and \mathcal{A} -continuous functions in biminimal structure spaces.

¹ นิสิตระดับปริญญาโท, ^{2,3} ผู้ช่วยศาสตราจารย์ ภาควิชาคณิตศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยมหาสารคาม

¹ Student faculty of Science, Mahasarakham University, MahaSarakhm, Thailand, 44150.

^{2,3} Assistant professor, Department of Mathematics, Faculty of Science, Mahasarakham University, Mahasarakham 44150, Thailand.

* Corresponding Author : jeeranunt.k@msu.ac.thdu

Preliminaries

In this section, we will give some definitions and notations, deal with some preliminaries and some useful results that will be duplicated in later sections.

Definition 2.1¹⁰ Let (X, τ) be a topological space and $M \subseteq X$. Then M is called an *A*-set if $M = U \cap B$ when U is open and B is regular closed in X .

The family of all *A*-sets in a topological space (X, τ) is denoted by $\mathcal{A}(X, \tau)$.

Definition 2.2¹⁷ Let X be a nonempty set and $P(X)$ be the power set of X . A subfamily m_x of $P(X)$ is called a minimal structure (briefly an *m*-structure) on X if $\emptyset \in m_x$ and $X \in m_x$.

The pair (X, m_x) , we denote a nonempty set X with an *m*-structure m_x on X and it is called a minimal structure space (briefly an *m*-space). Each member of m_x is said to be m_x -open and the complement of an m_x -open set is said to be m_x -closed.

Definition 2.3¹⁷ Let X be a nonempty set and m_x an *m*-structure on X . For a subset A of X the m_x -interior of A and the m_x -closure of A with respect to m_x are defined as follows:

$$m_x \text{Int}(A) = \cup \{ U : U \subseteq A, U \in m_x \},$$

$$m_x \text{Cl}(A) = \cap \{ F : A \subseteq F, X \setminus F \in m_x \}.$$

Lemma 2.4¹⁴ Let X be a nonempty set and m_x an *m*-structure on X . For any subsets A and B of X , the following properties hold:

- (1) $m_x \text{Cl}(X \setminus A) = X \setminus m_x \text{Int}(A)$ and $m_x \text{Int}(X \setminus A) = X \setminus m_x \text{Cl}(A)$,
- (2) If $(X \setminus A) \in m_x$, then $m_x \text{Cl}(A) = A$ and if $A \in m_x$, then $m_x \text{Int}(A) = A$,
- (3) $m_x \text{Cl}(\emptyset) = \emptyset$, $m_x \text{Cl}(X) = X$, $m_x \text{Int}(\emptyset) = \emptyset$ and $m_x \text{Int}(X) = X$,
- (4) If $A \subseteq B$, then $m_x \text{Cl}(A) \subseteq m_x \text{Cl}(B)$ and $m_x \text{Int}(A) \subseteq m_x \text{Int}(B)$,
- (5) $A \subseteq m_x \text{Cl}(A)$ and $m_x \text{Int}(A) \subseteq A$,
- (6) $m_x \text{Cl}(m_x \text{Cl}(A)) = m_x \text{Cl}(A)$ and $m_x \text{Int}(m_x \text{Int}(A)) = m_x \text{Int}(A)$,
- (7) $m_x \text{Int}(A \cap B) = m_x \text{Int}(A) \cap m_x \text{Int}(B)$ and $m_x \text{Int}(A) \cup m_x \text{Int}(B) \subseteq m_x \text{Int}(A \cup B)$,
- (8) $m_x \text{Cl}(A \cup B) = m_x \text{Cl}(A) \cup m_x \text{Cl}(B)$ and $m_x \text{Cl}(A \cap B) \subseteq m_x \text{Cl}(A) \cap m_x \text{Cl}(B)$.

Definition 2.5¹³ An *m*-structure m_x on a nonempty set A is said to have property \mathfrak{B} if the union of any family of subsets belonging to m_x belongs to m_x .

Lemma 2.6¹⁷ Let X be a nonempty set and m_x is an *m*-structure on X satisfying property \mathfrak{B} .

For $A \subseteq X$ the following properties hold:

- (1) $A \in m_x$ if and only if $m_x \text{Int}(A) = A$,
- (2) A is m_x -closed if and only if $m_x \text{Cl}(A) = A$,
- (3) $m_x \text{Int}(A)$ is m_x -open and $m_x \text{Cl}(A)$ is m_x -closed.

Definition 2.7³ Let (X, m_x) be an *m*-space and $R \subseteq X$. Then R is called m_x -regular closed if and only if $R = m_x \text{Cl}(m_x \text{Int}(R))$.

The family of all m_x -regular closed sets in an m_x -space (X, m_x) is denoted by $RC(X, m_x)$

Definition 2.8¹⁹ A subset A of an *m*-space (X, m_x) is called an *m*-preopen set if $A \subseteq m_x \text{Int}(m_x \text{Cl}(A))$ and an m_x -preclosed set if $m_x \text{Cl}(m_x \text{Int}(A)) \subseteq A$.

The family of all m_x -preopen sets in an *m*-space (X, m_x) is denoted by $PO(X, m_x)$ and m_x -preclosed sets in an *m*-space (X, m_x) is denoted by $PC(X, m_x)$

Definition 2.9¹⁹ Let (X, m_x) be an *m*-space and $A \subseteq X$, the m_x -preclosure of A is denoted by $m_x \text{pcl}(A)$ is defined as the intersection of all m_x -preclosed of (X, m_x) containing A .

Proposition 2.10¹⁹ Let (X, m_x) be an *m*-space and $A, B \subseteq X$. If $A \subseteq B$ then $m_x \text{pcl}(A) \subseteq m_x \text{pcl}(B)$.

Proposition 2.11¹⁹ Let (X, m_x) be an *m*-space and $A \subseteq X$. If m_x satisfies the property \mathfrak{B} . Then $m_x \text{pcl}(A) = A \cup m_x \text{Cl}(m_x \text{Int}(A))$.

Definition 2.12² Let A be a nonempty set and m_x^1, m_x^2 be *m*-structures on X . A triple (X, m_x^1, m_x^2) is called a bi-minimal structure space (briefly *bim*-space).

Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. The m_x -closure and m_x -interior of A with respect to m_x^i are denoted by $m_x^i \text{Cl}(A)$ and $m_x^i \text{Int}(A)$ respectively, for $i, j = 1, 2$.

Each member of m_x^i is said to be an m_x^i -open set and the complement of an open set is said to be m_x^i -closed, for $i, j = 1, 2$.

Definition 2.13⁴ A subset A of biminimal structure spaces (X, m_x^1, m_x^2) is said to be

- (1) $(i, j)m_x$ -regular open if $A = m_x^i \text{Int}(m_x^j \text{Cl}(A))$, where $i, j = 1$ or 2 and $i \neq j$,
- (2) $(i, j)m_x$ -semi-open if $A \subseteq m_x^i \text{Cl}(m_x^j \text{Int}(A))$ where $i, j = 1$ or 2 and $i \neq j$,
- (3) $(i, j)m_x$ -preopen if $A \subseteq m_x^i \text{Int}(m_x^j \text{Cl}(A))$, where $i, j = 1$ or 2 and $i \neq j$.

The complement of an $(i, j)m_x$ -regular open (resp. $(i, j)m_x$ -semi-open, $(i, j)m_x$ -preopen) set is called $(i, j)m_x$ -regular closed (resp. $(i, j)m_x$ -semi-closed, $(i, j)m_x$ -preclosed).

Lemma 2.14⁴ Let (X, m_x^1, m_x^2) be a biminimal structure space and A be a subset of X . Then

- (1) A is $(i, j)m_x$ -regular closed if and only if $A = m_x^i \text{Cl}(m_x^j \text{Int}(A))$,
- (2) A is $(i, j)m_x$ -semi-closed if and only if $m_x^i \text{Int}(m_x^j \text{Cl}(A)) \subseteq A$,
- (3) A is $(i, j)m_x$ -preclosed if and only if $m_x^i \text{Cl}(m_x^j \text{Int}(A)) \subseteq A$.

Definition 2.15 [4] Let (X, m_x^1, m_x^2) and (Y, m_y^1, m_y^2) be biminimal structure space. A function $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$ is said to be (i, j) - M -continuous at a point $x \in X$ and each $V \in m_y^i$ containing $f(x)$, there exists $U \in m_x^j$ containing x such that $f(U) \subseteq V$, where $i, j = 1$ or 2 and $i \neq j$.

A function $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$ is said to be (i, j) - M -continuous if it has this property at each point $x \in X$.

Theorem 2.16⁴ For a function $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$, the following properties are equivalent:

- (1) f is (i, j) - M -continuous;
- (2) $f^{-1}(V) = m_x^j \text{Int}(f^{-1}(V))$ for every $V \in m_y^i$
- (3) $f(m_x^i \text{Cl}(A)) \subseteq m_y^i \text{Cl}(f(A))$ for every subset A of X ;
- (4) $m_x^i \text{Cl}(f^{-1}(B)) \subseteq f^{-1}(m_y^i \text{Cl}(B))$ for every subset B of Y ;
- (5) $f^{-1}(m_y^i \text{Int}(B)) \subseteq m_x^j \text{Int}(f^{-1}(B))$ for every subset B of Y ;
- (6) $m_x^j \text{Cl}(f^{-1}(F)) = f^{-1}(F)$ for every m_y^i -closed set F of Y .

Results and Discussion

\mathcal{A} -sets in minimal structure space

In this section, we introduce the concept of \mathcal{A} -sets in minimal structure spaces.

Definition 3.1.1 Let (X, m_x) be an m -space. A subset M of A is said to be an m_x - \mathcal{A} -set if there exist G and R such that $M = G \cap R$ when G is m_x -open and R is m_x -regular closed.

The family of all m_x - \mathcal{A} -set in an m -space (X, m_x) is denoted by $\mathcal{A}(X, m_x)$.

Example 3.1.2 Let $X = \{1, 2, 3\}$. Define an m -structure m_x on X as follows : $m_x = \{\emptyset, \{2\}, \{1, 2\}, \{1, 3\}, X\}$. Then $RC(X, m_x) = \{\emptyset, \{2\}, \{1, 3\}, X\}$ and $\mathcal{A}(X, m_x) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{1, 3\}, X\}$.

Definition 3.1.3 Let (X, m_x) be an m -space and $A \subseteq X$, then A is said to be an m_x - \dagger -set if $m_x \text{Int}(A) = m_x \text{Int}(m_x \text{Cl}(A))$.

The family of all m_x - \dagger -set in an m -space (X, m_x) is denoted by $\dagger(X, m_x)$.

Example 3.1.4 Let $X = \{1, 2, 3\}$ and define $m_x = \{\emptyset, \{1\}, \{2\}, \{1, 3\}, \{2, 3\}, X\}$ be an m -structure on X . It follows that $\dagger(X, m_x) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 3\}, \{2, 3\}, X\}$.

Proposition 3.1.5 Let (X, m_x) be an m -space and $R \subseteq X$. If R is m_x -regular closed then R is an m_x - \dagger -set.

Proof. Let R be an m_x -regular closed. Then $R = m_x \text{Cl}(m_x \text{Int}(R))$. Consequently, $m_x \text{Cl}(R) = m_x \text{Cl}(m_x \text{Cl}(m_x \text{Int}(R)))$. Thus $m_x \text{Int}(m_x \text{Cl}(R)) = m_x \text{Int}(m_x \text{Cl}(m_x \text{Int}(R)))$. Hence $m_x \text{Int}(m_x \text{Cl}(R)) = m_x \text{Int}(R)$. Therefore, R is an m_x - \dagger -set.

\mathcal{A} -sets in biminimal structure space

In this section, we introduce the concept of \mathcal{A} -sets in biminimal structure spaces and study some fundamental properties of \mathcal{A} -sets in biminimal structure spaces and investigate some of their properties.

Definition 3.2.1 A subset A of a biminimal structure space (X, m_x^1, m_x^2) is said to be $(i, j)m_x$ -locally closed if there exist G and F such that $A = G \cap F$ when G is an m_x^i -open set and F is an m_x^j -closed set, where $i, j = 1$ or 2 and $i \neq j$.

The family of all $(i, j)m_x$ -locally closed sets in biminimal structure spaces (X, m_x^1, m_x^2) is denoted by $(i, j)m_x\text{-}\mathcal{LC}(X, m_x^1, m_x^2)$, where $i, j = 1$ or 2 and $i \neq j$.

Example 3.2.2 Let $X = \{a, b, c\}$. Define m -structures m_x^1 and m_x^2 on X as follows : $m_x^1 = \{\emptyset, \{b, c\}, X\}$ and $m_x^2 = \{\emptyset, \{c\}, X\}$. Thus $(1, 2)\text{-}\mathcal{LC}(X, m_x^1, m_x^2) = \{\emptyset, \{b\}, \{b, c\}, \{a, b\}, X\}$.

Lemma 3.2.3 Let S be a subset of a biminimal structure space (X, m_x^1, m_x^2) and let $i, j = 1$ or 2 and $i \neq j$. If S is an $(i, j)m_x$ -locally closed set then there exists an m_x^i -open set U such that $S = U \cap m_x^j Cl(S)$

Proof. Let S be an $(i, j)m_x$ -locally closed set. Then there exist U and F such that $S = U \cap F$ where U is m_x^i -open and F is m_x^j -closed. Since $S = U \cap F$, $S \subseteq F$. Thus $m_x^j Cl(S) \subseteq m_x^j Cl(F)$. Since F is m_x^j -closed, $m_x^j Cl(S) \subseteq F$. Then $U \cap m_x^j Cl(S) \subseteq U \cap F = S$. Since $S \subseteq U$ and $S \subseteq m_x^j Cl(S)$. Then $S \subseteq U \cap m_x^j Cl(S)$. Therefore, there exists an m_x^i -open set U such that $S = U \cap m_x^j Cl(S)$.

The converse of Lemma 3.2.3 is true if m_x^j has property \mathfrak{B} as a following proposition.

Proposition 3.2.4. Let S be a subset of a biminimal structure space (X, m_x^1, m_x^2) and let m_x^j has property \mathfrak{B} , where $i, j = 1$ or 2 and $i \neq j$. Then S is an $(i, j)m_x$ -locally closed set iff there exists an m_x^i -open set U such that $S = U \cap m_x^j Cl(S)$.

Proof. (\Rightarrow) By Lemma 3.2.3.

(\Leftarrow) Let $S = U \cap m_x^j Cl(S)$, for some $U \in m_x^i$. Since m_x^j has property \mathfrak{B} , $m_x^j Cl(S)$ is closed in (X, m_x^j) . Thus S is an $(i, j)m_x$ -locally closed.

Definition 3.2.5. Let (X, m_x^1, m_x^2) be a biminimal structure space. A subset M of X is said to be an $(i, j)m_x$ - \mathcal{A} -set if there exist G and R , such that $M = G \cap R$ when $G \in m_x^i$ and R is m_x^j -regular closed, where $i, j = 1$ or 2 and $i \neq j$.

The family of all $(i, j)m_x$ - \mathcal{A} -sets in a biminimal structure space (X, m_x^1, m_x^2) is denoted by $(i, j)\text{-}\mathcal{A}(X, m_x^1, m_x^2)$ where $i, j = 1$ or 2 and $i \neq j$.

Example 3.2.6. Let $X = \{1, 2, 3\}$. Define m -structures m_x^1 and m_x^2 on X as follows : $m_x^1 = \{\emptyset, \{1, 2\}, \{1, 3\}, X\}$ and $m_x^2 = \{\emptyset, \{2\}, \{1, 2\}, X\}$ which are m -structures on X . It follows that $RC(X, m_x^2) = \{\emptyset, X\}$. Thus $(1, 2)\text{-}\mathcal{A}(X, m_x^1, m_x^2) = \{\emptyset, \{1, 2\}, \{1, 3\}, X\}$.

Lemma 3.2.7. Let (X, m_x^1, m_x^2) be a biminimal structure space m_x^j has property \mathfrak{B} . If a subset M of X is an $(i, j)m_x$ - \mathcal{A} -set, then M is $(i, j)m_x$ -locally closed, where $i, j = 1$ or 2 and $i \neq j$.

Proof. Let M is an $(i, j)m_x$ - \mathcal{A} -set. Then there exist G and R such that $M = G \cap R$ where $G \in m_x^i$ -open and R is m_x^j -regular closed. Since R is m_x^j -regular closed, $R = m_x^j Cl(m_x^j Int(R))$. But m_x^j has property \mathfrak{B} , then $m_x^j Cl(m_x^j Int(R))$ is closed. Hence R is m_x^j -closed. It follows that M is an $(i, j)m_x$ -locally closed.

Proposition 3.2.8 Let (X, m_x^1, m_x^2) be a biminimal structure space and $m_x^j \subseteq m_x^i$ has the property \mathfrak{B} . If a subset M of X is both $(i, j)m_x$ -semi-open and $(i, j)m_x$ -locally closed, then M is an $(i, j)m_x$ - \mathcal{A} -sets, where $i, j = 1$ or 2 and $i \neq j$.

Proof. Let M be both $(i, j)m_x$ -semi-open and $(i, j)m_x$ -locally closed. It follows that $M \subseteq m_x^j Cl(m_x^j Int(M))$ such that $M = U \cap m_x^j Cl(M)$. Since $m_x^j Cl(M) \subseteq m_x^j Cl(m_x^j Int(M))$. But $m_x^j Cl(m_x^j Int(M)) \subseteq m_x^j Cl(M)$, hence $m_x^j Cl(M) = m_x^j Cl(m_x^j Int(M))$ and $m_x^j Cl(m_x^j Int(M))$ is m_x^j -regular closed. Consequently $m_x^j Cl(M)$ is m_x^j -regular closed. Therefore, M is an $(i, j)m_x$ - \mathcal{A} -set.

Definition 3.2.9 Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. Then A is said to be an $(i, j)m_x$ - \mathcal{t} -set if $m_x^j Int(A) = m_x^j Int(m_x^j Cl(A))$, where $i, j = 1$ or 2 and $i \neq j$.

The family of all $(i, j)m_x$ - \mathcal{t} -sets in a biminimal structure spaces (X, m_x^1, m_x^2) is denoted by $(i, j)\text{-}\mathcal{t}(X, m_x^1, m_x^2)$ for $i, j = 1$ or 2 and $i \neq j$.

Example 3.2.10 Let $X = \{1, 2, 3\}$. Define m -structures m_x^1 and m_x^2 on X as follows : $m_x^1 = \{\emptyset, \{1\}, \{3\}, \{2, 3\}, X\}$ and $m_x^2 = \{\emptyset, \{1\}, \{1, 2\}, X\}$.

Thus $(1, 2)\text{-}\mathcal{t}(X, m_x^1, m_x^2) = \{\emptyset, \{3\}, \{2, 3\}, X\}$.

Theorem 3.2.11 Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. Then A is an $(i, j)m_x$ - \mathcal{t} -set if and only if A is an $(i, j)m_x$ -semi-closed, where $i, j = 1$ or 2 and $i \neq j$.

Proof. (\Rightarrow) Let A be an $(i, j)m_x$ - \mathcal{t} -set. Then $m_x^j Int(A) = m_x^j Int(m_x^j Cl(A))$. Thus $m_x^j Int(m_x^j Cl(A)) \subseteq A$. Hence A is an $(i, j)m_x$ -semi-closed.

(\Leftarrow) Let A be an $(i, j)m_x$ -semi-closed. Then $m_x^j Int(m_x^j Cl(A)) \subseteq A$. Thus $m_x^j Int(m_x^j Int(m_x^j Cl(A))) \subseteq m_x^j Int(A)$. Hence $m_x^j Int(m_x^j Cl(A)) \subseteq m_x^j Int(A)$. Since $m_x^j Int(A) \subseteq m_x^j Int(m_x^j Cl(A))$. Thus $m_x^j Int(A) = m_x^j Int(m_x^j Cl(A))$. Hence A is an $(i, j)m_x$ - \mathcal{t} -set.

Definition 3.2.12 Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. Then A is said to be an $(i, j)m_x$ - \mathcal{B} -set if $A = U \cap T$, when U is an m_x^i -open set and T is an m_x^j - \mathcal{t} -set, where $i, j = 1$ or 2 and $i \neq j$.

The family of all $(i, j)m_x$ - \mathcal{B} -sets in a biminimal structure space (X, m_x^1, m_x^2) is denoted by $(i, j)\text{-}\mathcal{B}(X, m_x^1, m_x^2)$, where $i, j = 1$ or 2 and $i \neq j$.

Example 3.2.13 Let $X = \{1, 2, 3\}$. Define m -structures m_x^1 and m_x^2 on X as follows : $m_x^1 = \{\emptyset, \{1\}, \{2\}, \{2, 3\}, X\}$ and $m_x^2 = \{\emptyset, \{1\}, \{3\}, \{2, 3\}, X\}$. Then $\{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{2, 3\}, X\}$ are m_x^2 - \mathcal{t} -sets. Therefore, $(1, 2)\text{-}\mathcal{B}(X, m_x^1, m_x^2) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{2, 3\}, X\}$.

Theorem 3.2.14 Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. If A is an $(i, j)m_x$ - \mathcal{A} -set, then A is an $(i, j)m_x$ - \mathcal{B} -set for all $i, j = 1$ or 2 and $i \neq j$.

Proof. Let A be an $(i, j)m_x$ - \mathcal{A} -set. Then there exist G and R such that $A = G \cap R$ where G is m_x^i -open in (X, m_x^i) and R is an m_x^j -regular closed. By Proposition 3.1.5, R is an m_x^j - \mathcal{t} -set. Hence A is an $(i, j)m_x$ - \mathcal{B} -set.

Proposition 3.2.15 Let (X, m_x^1, m_x^2) be a biminimal structure space and M be a subset of X . If M is an $(i, j)m_x$ -locally closed set, then it is also an $(i, j)m_x$ - \mathcal{B} -set, where $i, j = 1$ or 2 and $i \neq j$.

Proof. Let M be $(i, j)m_x$ -locally closed set. Then there exist U and B such that $M = U \cap B$ where U is m_x^i -open and B is m_x^j -closed. Since B is m_x^j -closed, $B = m_x^jCl(B)$. Thus $m_x^jInt(B) = m_x^jInt(m_x^jCl(B))$. Hence B is an m_x^j - \mathcal{t} -set. Thus M is an $(i, j)m_x$ - \mathcal{B} -set.

Definition 3.2.16 Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. Then A is said to be an $(i, j)m_x$ - \mathcal{C} -set if $A = U \cap B$, when U is an m_x^i -open and B is m_x^j -preclosed, where $i, j = 1$ or 2 and $i \neq j$.

The family of all $(i, j)m_x$ - \mathcal{C} -sets in a biminimal structure space (X, m_x^1, m_x^2) is denoted by $(i, j)\text{-}\mathcal{C}(X, m_x^1, m_x^2)$, where $i, j = 1$ or 2 and $i \neq j$.

Example 3.2.17 Let $X = \{1, 2, 3\}$. Define m -structures m_x^1 and m_x^2 on X as follows : $m_x^1 = \{\emptyset, \{1\}, \{2\}, \{2, 3\}, X\}$ and $m_x^2 = \{\emptyset, \{1\}, \{3\}, \{2, 3\}, X\}$. Thus $\emptyset, \{1\}, \{2\}, \{1, 2\}, \{2, 3\}, X$ are preclosed in (X, m_x^2) . Therefore, $(1, 2)\text{-}\mathcal{C}(X, m_x^1, m_x^2) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{2, 3\}, X\}$.

Theorem 3.2.18. Let (X, m_x^1, m_x^2) be a biminimal structure space and $A \subseteq X$. If A is an $(i, j)m_x$ - \mathcal{A} -set, then A is an $(i, j)m_x$ - \mathcal{C} -set for all $i, j = 1$ or 2 and $i \neq j$.

Proof Let A be an $(i, j)m_x$ - \mathcal{A} -set. Then there exist G and R such that $A = G \cap R$ where G is m_x^i -open and R is an m_x^j -regular closed. Since $R = m_x^jCl(m_x^jInt(R))$, thus

$m_x^jCl(m_x^jInt(R)) \subseteq R$. Hence R is an m_x^j -preclosed. Therefore, A is an $(i, j)m_x$ - \mathcal{C} -set.

Proposition 3.2.19 Let (X, m_x^1, m_x^2) be a biminimal structure space and M be a subset of X . If M is an $(i, j)m_x$ -locally closed set, then it is also an $(i, j)m_x$ - \mathcal{C} -set, where $i, j = 1$ or 2 and $i \neq j$.

Proof. Let M be an $(i, j)m_x$ -locally closed set. Then there exist U and B such that $M = U \cap B$ where U is m_x^i -open in (X, m_x^i) and B is m_x^j -closed.

It follows that $m_x^jCl(m_x^jInt(B)) \subseteq m_x^jCl(B) = B$. Then B is an m_x^j -preclosed. Hence M is an $(i, j)m_x$ - \mathcal{C} -set.

Proposition 3.2.20 Let A be a subset of a biminimal structure space (X, m_x^1, m_x^2) and m_x^j has the property \mathfrak{B} . Then A is an $(i, j)m_x$ - \mathcal{C} -set iff $A = U \cap m_x^jpcl(A)$ for some $U \in m_x^i$, where $i, j = 1$ or 2 and $i \neq j$.

Proof. (\Rightarrow) Let A be $(i, j)m_x$ - \mathcal{C} -set. Then there exist U and B such that $A = U \cap B$ where U is m_x^i -open and B is m_x^j -preclosed. From $A \subseteq B$, $m_x^jpcl(A) \subseteq m_x^jpcl(B)$ by Proposition 2.11, $m_x^jpcl(B) = B \cup m_x^jCl(m_x^jInt(B))$. As B is m_x^j -preclosed, $m_x^jCl(m_x^jInt(B)) \subseteq B$. Hence $m_x^jpcl(B) = B$. Thus $m_x^jpcl(A) \subseteq B$. Therefore $A = U \cap m_x^jpcl(A)$.

(\Leftarrow) Let $A = U \cap m_x^jpcl(A)$ for some $U \in m_x^i$. Since $m_x^jpcl(A)$ is an m_x^j -preclosed. Therefore, A is an $(i, j)m_x$ - \mathcal{C} -set.

Proposition 3.2.21 Let A be a subset of a biminimal structure space (X, m_x^1, m_x^2) and m_x^j has the property \mathfrak{B} . Then $A = U \cap m_x^jCl(m_x^jInt(A))$ for some $U \in m_x^i$ if and only if A is an m_x^i -semi-open and $(i, j)m_x$ - \mathcal{C} -set, where $i, j = 1$ or 2 and $i \neq j$.

Proof. (\Rightarrow) Let $A = U \cap m_x^jCl(m_x^jInt(A))$ for some $U \in m_x^i$. Then $A \subseteq m_x^jCl(m_x^jInt(A))$. Thus A is an m_x^i -semi-open. By Lemma 2.6, $m_x^jCl(m_x^jInt(A))$ is m_x^j -closed. Since $m_x^jInt(m_x^jCl(m_x^jInt(A))) \subseteq m_x^jCl(m_x^jInt(A))$, $m_x^jCl(m_x^jInt(m_x^jCl(m_x^jInt(A)))) \subseteq m_x^jCl(m_x^jInt(A))$. Hence $m_x^jCl(m_x^jInt(A))$ is m_x^j -preclosed. Then A is an $(i, j)m_x$ - \mathcal{C} -set.

(\Leftarrow) Let A be an m_x^i -semi-open and $(i, j)m_x$ - \mathcal{C} -set. By Proposition 3.2.20, $A = U \cap m_x^jpcl(A)$ for some $U \in m_x^i$. Since A is m_x^i -semi-open. Then $A \subseteq m_x^jCl(m_x^jInt(A))$. Since m_x^j has the property \mathfrak{B} and by Proposition 2.11, $m_x^jpcl(A) = A \cup m_x^jCl(m_x^jInt(A))$. Thus $m_x^jpcl(A) = m_x^jCl(m_x^jInt(A))$. Hence $A = U \cap m_x^jCl(m_x^jInt(A))$

for some $U \in m_x^i$.

Theorem 3.2.22 Let A be a subset of a biminimal structure space (X, m_x^1, m_x^2) and m_x^i has the property \mathfrak{B} . If a subset M of X is an m_x^i -semi-open and $(i, j)m_x^i$ - \mathcal{C} -set, then it is an $(i, j)m_x^i$ - \mathcal{A} -set.

Proof. Let M be an m_x^i -semi-open and $(i, j)m_x^i$ - \mathcal{C} -set. By Proposition 3.2.21, then $M = U \cap m_x^i Cl(m_x^i Int(M))$, for some $U \in m_x^i$. Since $m_x^i Cl(m_x^i Int(M))$ is m_x^i -regular closed. Therefore, M is an $(i, j)m_x^i$ - \mathcal{A} -set.

Definition 3.2.23 Let (X, m_x^1, m_x^2) and (Y, m_y^1, m_y^2) be biminimal structure spaces.

A function $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$ is said to be

- (1) (i, j) -semi-continuous if $f^{-1}(V) \in (i, j)$ - $SO(X, m_x^1, m_x^2)$ for all $V \in m_y^i$.
- (1) (i, j) - \mathcal{LC} -continuous if $f^{-1}(V) \in (i, j)$ - $\mathcal{LC}(X, m_x^1, m_x^2)$ for all $V \in m_y^i$.
- (1) (i, j) - \mathcal{A} -continuous if $f^{-1}(V) \in (i, j)$ - $\mathcal{A}(X, m_x^1, m_x^2)$ for all $V \in m_y^i$.
- (1) (i, j) - \mathcal{C} -continuous if $f^{-1}(V) \in (i, j)$ - $\mathcal{C}(X, m_x^1, m_x^2)$ for all $V \in m_y^i$.

Proposition 3.2.24 Let (X, m_x^1, m_x^2) and (Y, m_y^1, m_y^2) be biminimal structure spaces. A function $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$ be a mapping. If f is (i, j) - \mathcal{A} -continuous then f is (i, j) - \mathcal{LC} -continuous.

Proof. Let f be (i, j) - \mathcal{A} -continuous and $V \in m_y^i$. Then $f^{-1}(V) \in (i, j)$ - $\mathcal{A}(X, m_x^1, m_x^2)$. By Lemma 3.2.7, we have $f^{-1}(V) \in (i, j)$ - $\mathcal{LC}(X, m_x^1, m_x^2)$. Hence f is (i, j) - \mathcal{LC} -continuous.

Theorem 3.2.25 Let (X, m_x^1, m_x^2) and (Y, m_y^1, m_y^2) be biminimal structure spaces and m_x^i has the property \mathfrak{B} . If a mapping $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$ is (i, j) -semi-continuous and (i, j) - \mathcal{LC} -continuous then f is (i, j) - \mathcal{A} -continuous.


Proof. Let f be an (i, j) -semi-continuous and (i, j) - \mathcal{LC} -continuous and $V \in m_y^i$. Then $f^{-1}(V) \in (i, j)$ - $SO(X, m_x^1, m_x^2)$ and $f^{-1}(V) \in (i, j)$ - $\mathcal{LC}(X, m_x^1, m_x^2)$. By Proposition 3.2.8, thus $f^{-1}(V) \in (i, j)$ - $\mathcal{A}(X, m_x^1, m_x^2)$. Therefore, f is (i, j) - \mathcal{A} -continuous.

Theorem 3.2.26 Let (X, m_x^1, m_x^2) and (Y, m_y^1, m_y^2) be biminimal structure spaces. If a mapping $f : (X, m_x^1, m_x^2) \rightarrow (Y, m_y^1, m_y^2)$ is (i, j) -semi-continuous and (i, j) - \mathcal{C} -continuous then f is (i, j) - \mathcal{A} -continuous.

Proof. Let f be an (i, j) -semi-continuous and (i, j) - \mathcal{C} -continuous and $V \in m_y^i$. Then $f^{-1}(V) \in (i, j)$ - $SO(X, m_x^1, m_x^2)$ and $f^{-1}(V) \in (i, j)$ - $\mathcal{C}(X, m_x^1, m_x^2)$. By Theorem 3.2.22, thus $f^{-1}(V) \in (i, j)$ - $\mathcal{A}(X, m_x^1, m_x^2)$. Therefore, f is (i, j) - \mathcal{A} -continuous.

Conclusion

In this paper, we introduced the concept of \mathcal{A} -sets in biminimal structure spaces. We also studied some properties of \mathcal{A} -sets and \mathcal{A} -continuous function on the space. The following implications hold for a biminimal structure spaces. These implications are not reversible.



Acknowledgement

The authors would like to thank the referees for helpful comments and suggestions on the manuscript.

References

1. Andrijevic D. *Semi-preopen sets*. (1986), Math. Vesnik., 38, 24-32.
2. Boonpok C. *Almost and weakly M-continuous functions in m-spaces*. Far East J.Math.Sci, 43 (2010), 29-40
3. Boonpok C. *Biminimal structure space*. (2010), Int. Math. Forum, 5(15), 703-707
4. Boonpok C. *M-continuous functions on biminimal structure space*. Far East J.Math.Sci, 43 (2010), 41-58
5. Bourbaki N. *General Topology. Part 1*. (1996), Massachusetts, Addison-Wesley.
6. Császár Á. *connected sets*. Acta Math. Hungar. (2003), 101(4): 273-279.
7. Dugundji J. *Topology*. (1972), Boston, Allyn and Bacon.
8. Erguang Y and Pengfei Y. *On decomposition of A-continuity*. (2006), Acta Math. Hungar., 110, 309-313.
9. Ganster M. and Reilly, I. L. *A decomposition of continuity*. (1990). Acta Math.Hungar., 56, 299-301.

10. Kelly J.C *Bitopological space*. (1969), Pro.London Math.Soc.,39130,71-79
11. Keun Min W. *m-open set and M-continuous functions*. (2010), Commun. Korean. Math. Soc. 25, No.2, pp 251-256
12. Levine N. *Semi-open sets and semi-continuity in topological spaces*. (1963), Amer. Math. Monthly, 70, 36-41.
13. Mashhour A, Abd El-Monsef M, and El-Deeb S. *On pre-continuous and weak precontinuous mappings*. (1982), Proc. Math. and Phys. Soc. Egypt., 53, 47-53.
14. Maki H, Umehara J, Noiri T. *Every topological space is pre $T_{1/2}$* . Fac. Sci. Kochi Univ. ser. Math., pp 33-42
15. Maki H, Rao K.C, Nagoor Gani A. *On generalized semi-open and preopen sets*. Pure Appl. Math. Sci. 1999; 49: 17-29.
16. Nj_astad O. *On some classes of nearly open sets*. (1965), Pacific J. Math., 15, 961-970.
17. Noiri T and Popa V. *A generalized of some forms of g-irresolute function*. European J. of Pure and Appl. Math. 2(4)(2009), 473-493
18. Popa V and Noiri T. *On M-continuous function*.(1965), Anal.Univ. "Dunarea de Jos" Galati, Ser.Mat.Fiz.Mec. Teor.(2),18(23), 31-41
19. Rosas E. *Some new types of open and closed sets in minimal structures-II*. (2009), Int. Math. Forum, No.44, 2185-2198.
20. Suantai S. *Topology*. Department of Mathematics, Faculty of Science, Chiang Mai University.
21. Tong J. *A decomposition of continuity*. (1986), Acta Math. Hungar., 48, 11-15.
22. Tong J. *On decomposition of continuity in topological spaces*. (1989), Acta Math. Hungar., 48, 11-15.