

# สมบัติของตัวดำเนินการ $\mathcal{R}_m^a$ ในปริภูมิโครงสร้างเล็กสุดที่มีอุดมคติ

## Properties of $\mathcal{R}_m^a$ -operator in minimal structure space with an ideal

ยุทธพงศ์ มนต์พิพารมณ<sup>1</sup>, ดรุณี บุญชารี<sup>2\*</sup>, โชคชัย วิริยะพงษ์<sup>3</sup>,  
Yutthapong Manuttiparom<sup>1</sup>, Daruni Boonchari<sup>2\*</sup>, Chokchai Viriyapong<sup>3</sup>

Received: 19 October 2020 ; Revised: 20 December 2020 ; Accepted: 8 January 2021

### บทคัดย่อ

ในบทความนี้ ผู้วิจัยได้นำเสนอเซตเปิดแบบ  $\delta$ - $m$  เซตเปิดแบบ  $a$ - $m$  ฟังก์ชันแบบ  $\delta$ - $m$ -local ตัวดำเนินการแบบ  $R_m^a$  บนปริภูมิโครงสร้างเล็กสุดที่มีอุดมคติพร้อมทั้งศึกษาสมบัติของฟังก์ชัน และตัวดำเนินการนี้

**คำสำคัญ:** เซตเปิดแบบ  $\delta$ - $m$  เซตเปิดแบบ  $a$ - $m$  ฟังก์ชันแบบ  $\delta$ - $m$ -local ตัวดำเนินการแบบ  $R_m^a$  ปริภูมิโครงสร้างเล็กสุดที่มีอุดมคติ

### Abstract

In this article, the concepts of  $\delta$ - $m$ -open sets,  $a$ - $m$ -open sets in a minimal structure space with an ideal are introduced. In addition, we present an  $a$ - $m$ -local function and an  $R_m^a$ -operator in a minimal structure space with an ideal. We studied the properties of the function and this operator.

**Keywords:**  $\delta$ - $m$ -open sets,  $a$ - $m$ -open sets,  $\delta$ - $m$ -local functions,  $R_m^a$ -operator, a minimal structure space with an ideal.

---

<sup>1</sup> นิสิตปริญญาโท, คณะวิทยาศาสตร์ มหาวิทยาลัยมหาสารคาม อำเภอกันทรวิชัย จังหวัดมหาสารคาม 44150

<sup>2</sup> ผู้ช่วยศาสตราจารย์, คณะวิทยาศาสตร์ มหาวิทยาลัยมหาสารคาม อำเภอกันทรวิชัย จังหวัดมหาสารคาม 44150

<sup>3</sup> ผู้ช่วยศาสตราจารย์, คณะวิทยาศาสตร์ มหาวิทยาลัยมหาสารคาม อำเภอกันทรวิชัย จังหวัดมหาสารคาม 44150

<sup>1</sup> Master degree student, Mahasarakham University, Kantharawichai District, Maha Sarakham 44150, Thailand.

<sup>2</sup> Asst. Prof., Faculty of Science, Mahasarakham University, Kantharawichai District, Maha Sarakham 44150, Thailand.

<sup>3</sup> Asst. Prof., Faculty of Science, Mahasarakham University, Kantharawichai District, Maha Sarakham 44150, Thailand.

\* Corresponding author ; Daruni Boonchari, Faculty of Science, Mahasarakham University, Kantharawichai District, Maha Sarakham 44150, Thailand. daruni.b@msu.ac.th.

## Introduction

In 1945, Vaidyanathaswamy (1945) defined a local function in an ideal topological space and studied some properties of this function. In 1996, Maki, Umehara and Noiri (1996) defined a minimal structure and studied some properties of this structure. In 2014, Al-Omeri *et al.* (2014) defined an *a-local* function in an ideal topological space and also studied some properties of an *a-local* function. Later in 2016, Al-Omeri *et al.* (2016) defined an  $R_a$ -operator in an ideal topological space and studied some properties of this operator. In this article, we introduce the concepts of  $\delta$ -*m-open* sets and  $\delta$ -*m-closed* sets in a minimal structure space with an ideal and study some fundamental properties. Moreover, we introduce the notions of  $\delta$ -*m-local* functions and  $R_m^a$ -operators in minimal structure spaces, along with studying some properties related to an  $\delta$ -*m-local* function and an  $R_m^a$ -operator defined above.

## Preliminaries

Definition 2.1<sup>5</sup> Let  $X$  be a nonempty set and  $P(X)$  the power set of  $X$ . A subfamily  $m$  of  $P(X)$  is called a minimal structure (briefly *MS*) on  $X$  if  $\emptyset \in m$  and  $X \in m$ .

By  $(X, m)$  we denote a nonempty set  $X$  with a minimal structure  $m$  on  $X$  and it is called a minimal structure space. Each member of  $m$  is said to be *m-open* and the complement of *m-open* is said to be *m-closed*.

Definition 2.2 (Noiri & Popa, 2009) Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . The *m-closure* of  $A$ , denoted by  $CI_m(A)$  and the *m-interior* of  $A$ , denoted by  $Int_m(A)$ , are defined as follows ;

- 1)  $CI_m(A) = \bigcap \{F : A \subseteq F, X \setminus F \in m\}$ ,
- 2)  $Int_m(A) = \bigcup \{U : U \subseteq A, U \in m\}$ .

Lemma 2.3 (Maki & Gani, 1999) Let  $(X, m)$  be a minimal structure space and  $A, B \subseteq X$ , the following properties hold ;

- (1)  $CI_m(X \setminus A) = X \setminus Int_m(A)$  and  $Int_m(X \setminus A) = X \setminus CI_m(A)$ .
- (2) If  $X \setminus A \in m$ , then  $CI_m(A) = A$  and if  $A \in m$ , then  $Int_m(A) = A$ .
- (3)  $CI_m(\emptyset) = \emptyset$ ,  $CI_m(X) = X$ ,  $Int_m(\emptyset) = \emptyset$ , and  $Int_m(X) = X$ .

(4) If  $A \subseteq B$ , then  $CI_m(A) \subseteq CI_m(B)$  and  $Int_m(A) \subseteq Int_m(B)$ .

(5)  $A \subseteq CI_m(A)$  and  $Int_m(A) \subseteq A$ .

(6)  $CI_m(CI_m(A)) = CI_m(A)$  and  $Int_m(Int_m(A)) = Int_m(A)$ .

Lemma 2.4 (Maki & Gani, 1999) Let  $(X, m)$  be a minimal structure space and  $A \subseteq X, x \in X$ . Then  $x \in CI_m(A)$  if and only if  $U \cap A \neq \emptyset$  for every an *m-open* set  $U$  containing  $x$ .

Definition 2.5 (Rosas *et al.*, 2009) Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ .

(1)  $A$  is called *m-regular open* if  $A = Int_m(CI_m(A))$

(2)  $A$  is called *m-regular closed* if  $X \setminus A$  is *m-regular open*.

The family of all *m-regular open* sets of  $X$  is denoted by  $r(m)$  and the family of all *m-regular closed* sets of  $X$  is denoted by  $rc(m)$ .

Definition 2.6 (Ozbakir & Yildirim, 2009) An ideal  $\mathcal{I}$  on a minimal structure space  $(X, m)$  is a nonempty collection of subsets of  $X$  which satisfies the following properties ;

- (1)  $A \in \mathcal{I}$  and  $B \subseteq A$  implies  $B \in \mathcal{I}$  (heredity),
- (2)  $A \in \mathcal{I}$  and  $B \in \mathcal{I}$  implies  $A \cup B \in \mathcal{I}$  (finite additivity).

The set  $\mathcal{I}$  together with a minimal structure space  $(X, m)$  is called a minimal structure space with an ideal, denoted by  $(X, m, \mathcal{I})$ .

## Main Results

Definition 3.1 Let  $(X, m)$  be a minimal structure space. A subset  $A$  is said to be  $\delta$ -*m-open* if for each  $X \in A$  there exists an *m-regular open* set  $G$  such that  $X \in G \subseteq A$ . The complement of  $\delta$ -*m-open* set is called  $\delta$ -*m-closed*. The family of all  $\delta$ -*m-closed* sets of  $X$ , denoted by  $\delta C_m(X)$ .

Theorem 3.2 Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . The arbitrary union of  $\delta$ -*m-open* sets is a  $\delta$ -*m-open* set.

Proof Let  $B_\alpha$  be a  $\delta$ -*m-open* set for all  $\alpha \in J$  where  $J$  is an index set and let  $x \in \bigcup_{\alpha \in J} B_\alpha$ . There exists  $\beta \in J$  such that  $x \in B_\beta$ . Since  $B_\beta$  is  $\delta$ -*m-open*, there exists an *m-regular open* set  $G_\beta$  such that  $x \in G_\beta \subseteq B_\beta$ . Then  $x \in G_\beta \subseteq B_\beta \subseteq \bigcup_{\alpha \in J} B_\alpha$ . Therefore  $\bigcup_{\alpha \in J} B_\alpha$  is  $\delta$ -*m-open*.

**Definition 3.3** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . A point  $x \in X$  is called a  $\delta$ - $m$ -cluster point of  $A$  if  $U \cap A \neq \emptyset$  for each  $m$ -regular open set  $U$  containing  $x$ .

**Definition 3.4** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . The set of all  $\delta$ - $m$ -cluster points of  $A$  is called  $\delta$ - $m$ -closure of  $A$  and is denoted by  $C_{\delta m}(A)$  and the union  $m$ -regular open sets contained in  $A$  is called the  $\delta$ - $m$ -interior of  $A$ , denoted by  $I_{\delta m}(A)$ .

**Theorem 3.5** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . Then  $A$  is  $\delta$ - $m$ -open if and only if  $I_{\delta m}(A) = A$ .

**Proof** ( $\Rightarrow$ ) Suppose that  $A$  is  $\delta$ - $m$ -open. By definition of  $\delta$ - $m$ -interior,  $I_{\delta m}(A) = A$ . Let  $x \in A$ . Since  $A$  is  $\delta$ - $m$ -open, there exists an  $m$ -regular open set  $O$  such that  $x \in O \subseteq A$ . This implies that  $x \in I_{\delta m}(A)$ . Then  $A \subseteq I_{\delta m}(A)$ . Hence  $A = I_{\delta m}(A) = A$ . ( $\Leftarrow$ ) It follows from Theorem 3.2.

**Theorem 3.6** Let  $(X, m)$  be a minimal structure space and  $A, B \subseteq X$ . The following property hold ;

- (1) If  $A \subseteq B$ , then  $I_{\delta m}(A) \subseteq I_{\delta m}(B)$ ,
- (2) If  $A \subseteq B$ , then  $C_{\delta m}(A) \subseteq C_{\delta m}(B)$ .

**Proof** (1) Assume that  $A \subseteq B$  and  $x \in I_{\delta m}(A)$ . Then, there exists an  $m$ -regular open set  $G$  such that  $x \in G \subseteq A$ . Since  $A \subseteq B$ , we have  $x \in G \subseteq A \subseteq B$ . This implies that  $x \in I_{\delta m}(B)$ . Hence  $I_{\delta m}(A) \subseteq I_{\delta m}(B)$ .

(2) Let  $A \subseteq B$ . Assume that  $x \notin C_{\delta m}(B)$ . Then there exists an  $m$ -regular open set  $U$  containing  $x$  such that  $U \cap B = \emptyset$ . Since  $A \subseteq B$ , we have  $U \cap A \subseteq U \cap B = \emptyset$ . Thus  $x \notin C_{\delta m}(A)$ . Therefore  $C_{\delta m}(A) \subseteq C_{\delta m}(B)$ .

**Theorem 3.7** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . The following properties hold ;

- (1)  $C_{\delta m}(A) = X \setminus I_{\delta m}(X \setminus A)$ ,
- (2)  $I_{\delta m}(A) = X \setminus C_{\delta m}(X \setminus A)$ .

**Proof** (1) We will show that  $C_{\delta m}(A) = X \setminus I_{\delta m}(X \setminus A)$  by contrapositive. Assume that  $x \notin X \setminus I_{\delta m}(X \setminus A)$ . We get that  $x \in I_{\delta m}(X \setminus A)$ . So there exists an  $m$ -regular open set  $G$  such that  $x \in G \subseteq X \setminus A$ . Then  $G \cap A = \emptyset$  and  $x \notin C_{\delta m}(A)$ . Thus  $C_{\delta m}(A) \subseteq X \setminus I_{\delta m}(X \setminus A)$ .

Next, we show that  $X \setminus I_{\delta m}(X \setminus A) \subseteq C_{\delta m}(A)$  by contrapositive. Assume that  $x \notin C_{\delta m}(A)$ . Then  $x$  is not a  $\delta$ - $m$ -cluster point of  $A$ . There exists an  $m$ -regular open set  $G$  containing  $x$  such that  $G \cap A = \emptyset$ .

So  $x \in G \subseteq X \setminus A$  and we get that  $x \in I_{\delta m}(X \setminus A)$ . Hence  $x \notin X \setminus I_{\delta m}(X \setminus A)$ . Thus  $X \setminus I_{\delta m}(X \setminus A) \subseteq C_{\delta m}(A)$ .

(2) Since  $X \setminus A \subseteq X$ , we have  $C_{\delta m}(X \setminus A) = X \setminus I_{\delta m}(X \setminus (X \setminus A))$  by (1) and we get  $C_{\delta m}(X \setminus A) = X \setminus I_{\delta m}(A)$ . Therefore  $I_{\delta m}(X \setminus A) = X \setminus C_{\delta m}(X \setminus A)$ .

**Definition 3.8** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ .

- (1)  $A$  is called  $a$ - $m$ -open if  $A \subseteq \text{Int}_m(CI_m(I_{\delta m}(A)))$ . The family of all  $a$ - $m$ -open sets of  $X$  is denoted by  $\mathcal{M}^a$ .
- (2)  $A$  is called  $a$ - $m$ -closed if  $CI_{\delta m}(\text{Int}_m(C_{\delta m}(A))) \subseteq A$ .

**Theorem 3.9** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . Then  $A$  is  $a$ - $m$ -open if and only if  $X \setminus A$  is  $a$ - $m$ -closed.

**Proof** Assume that  $A$  is  $a$ - $m$ -open. Then  $A \subseteq \text{Int}_m(CI_m(I_{\delta m}(A)))$ . and  $X \setminus A \supseteq X \setminus (\text{Int}_m(CI_m(I_{\delta m}(A))))$ . By Lemma 2.3 and Theorem 3.7,  $X \setminus A \supseteq CI_m(\text{Int}_m(C_{\delta m}(X \setminus A)))$ . Therefore,  $X \setminus A$  is  $a$ - $m$ -closed.

Conversely, assume that  $X \setminus A$  is  $a$ - $m$ -closed. Then  $CI_m(\text{Int}_m(C_{\delta m}(X \setminus A))) \subseteq X \setminus A$  and  $X \setminus CI_m(\text{Int}_m(C_{\delta m}(X \setminus A))) \supseteq X \setminus (X \setminus A)$ . By Lemma 2.3 and Theorem 3.7,  $\text{Int}_m(CI_m(I_{\delta m}(A))) \supseteq A$ . Hence  $A$  is  $a$ - $m$ -open.

**Example 3.10** Let  $X = \{a, b, c, d\}$  with a minimal structure  $m = \{\emptyset, \{a, b\}, \{b, c\}, \{c, d\}, \{a, d\}, x\}$ . Then  $r(m) = \{\emptyset, \{a, b\}, \{a, d\}, \{b, c\}, \{c, d\}, x\}$ , and  $\delta O_m(x) = \{\emptyset, \{a, b\}, \{a, d\}, \{b, c\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}, x\}$ ,  $\mathcal{M}^a = \{\emptyset, \{a, b\}, \{a, d\}, \{b, c\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}, x\}$ . In this example  $\{a, b\}, \{a, d\} \in \mathcal{M}^a$  but  $\{a, b\} \cap \{a, d\} = \{a\} \notin \mathcal{M}^a$ , that means  $\mathcal{M}^a$  does not have the property that any finite intersection of  $a$ - $m$ -open sets is  $a$ - $m$ -open.

**Definition 3.11** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ . The  $a$ - $m$ -closure of  $A$ , denoted by  $aC_m(A)$  and the  $a$ - $m$ -interior of  $A$ , denoted by  $aI_m(A)$ , are defined as follows ;

- (1)  $aC_m(A) = \bigcap \{F : X \setminus F \in \mathcal{M}^a \text{ and } A \subseteq F\}$ ,
- (2)  $aI_m(A) = \bigcup \{U : U \in \mathcal{M}^a \text{ and } U \subseteq A\}$ .

**Theorem 3.12** Let  $(X, m)$  be a minimal structure space and  $A \subseteq X$ ,  $x \in X$ , Then  $x \in aC_m(A)$  if and only if  $U \cap A \neq \emptyset$  for every  $a$ - $m$ -open set  $U$  containing  $x$ .

**Proof** ( $\Rightarrow$ ) Suppose that there exists an  $a$ - $m$ -open set  $U$  containing  $x$  such that  $U \cap A = \emptyset$ . So  $A \subseteq X \setminus U$  and  $X \setminus U$  is  $a$ - $m$ -closed. Since  $aC_m(A)$  is the intersection of

all  $a$ - $m$ -closed sets containing  $A$ ,  $aC_m(A) \subseteq X \setminus U$ . Since  $x \notin X \setminus U$ , we have  $x \notin aC_m(A)$ .

( $\Leftarrow$ ) Assume that  $x \notin aC_m(A)$ . Then there exists an  $a$ - $m$ -closed set  $F$  such that  $A \subseteq F$  and  $x \notin F$ . Choose  $U = X \setminus F$ . Then  $U$  is  $a$ - $m$ -open and  $x \in X \setminus F = U$ . Moreover,  $U \cap A \subseteq (X \setminus F) \cap F = \emptyset$ .

**Theorem 3.13** Let  $(X,m)$  be a minimal structure space and  $A, B \subseteq X$ . The following properties hold ;

- (1) If  $A \subseteq B$ , then  $aC_m(A) \subseteq aC_m(B)$ .
- (2) If  $A \subseteq B$ , then  $aI_m(A) \subseteq aI_m(B)$ .

**Proof** (1) Assume that  $A \subseteq B$  and  $x \notin aC_m(B)$ . Then there exists an  $a$ - $m$ -open set  $U$  containing  $x$  such that  $U \cap B = \emptyset$ . Since  $A \subseteq B$ ,  $U \cap A = \emptyset$ . Hence  $x \notin aC_m(A)$ .

(2) Let  $A \subseteq B$  and  $x \in aI_m(A)$ . Then there exists an  $a$ - $m$ -open set  $U$  such that  $x \in U \subseteq A$ . Since  $A \subseteq B$ ,  $x \in U \subseteq B$ . Therefore  $x \in aI_m(B)$ .

**Proposition 3.14** Let  $(X,m)$  be a minimal structure space. Then  $\emptyset \in \mathcal{M}^a$  and  $X \in \mathcal{M}^a$ .

**Proof** Since  $\emptyset \subseteq Int_m(CI_m(I_{\delta m}(\emptyset)))$ ,  $\emptyset$  is  $a$ - $m$ -open, and so  $\emptyset \in \mathcal{M}^a$ . Clearly  $X = Int_m(CI_m(X))$ , so  $X$  is an  $m$ -regular open. Then  $X$  is  $\delta$ - $m$ -open, that is  $I_{\delta m}(X) = X$ , and so  $X \subseteq Int_m(CI_m(I_{\delta m}(X)))$ . Therefore  $X \in \mathcal{M}^a$ .

**Theorem 3.15** Let  $(X,m)$  be a minimal structure space. Then the arbitrary union of elements of  $\mathcal{M}^a$  belongs to  $\mathcal{M}^a$ .

**Proof** Let  $V_\alpha$  be  $a$ - $m$ -open for all  $\alpha \in J$  and  $G = \bigcup_{\alpha \in J} V_\alpha$ . Then  $V_\alpha \subseteq Int_m(CI_m(I_{\delta m}(V_\alpha)))$  for all  $\alpha \in J$ . Since  $V_\alpha \subseteq G$ , it follows that  $I_{\delta m}(V_\alpha) \subseteq I_{\delta m}(G)$  and so  $CI_m(I_{\delta m}(V_\alpha)) \subseteq CI_m(I_{\delta m}(G))$ . Then  $Int_m(CI_m(I_{\delta m}(V_\alpha))) \subseteq Int_m(CI_m(I_{\delta m}(G)))$ . This implies that  $V_\alpha \subseteq Int_m(CI_m(I_{\delta m}(G)))$  for all  $\alpha \in J$ . Thus  $\bigcup_{\alpha \in J} V_\alpha \subseteq Int_m(CI_m(I_{\delta m}(G)))$ . Therefore  $G \subseteq Int_m(CI_m(I_{\delta m}(G)))$ .

**Corollary 3.16** Let  $(X,m)$  be a minimal structure space. Then the arbitrary intersection of  $a$ - $m$ -closed sets is an  $a$ - $m$ -closed set.

**Proof** Let  $G_\alpha$  be  $a$ - $m$ -closed for all  $\alpha \in J$ . Then  $X \setminus G_\alpha$  is  $a$ - $m$ -open and so  $\bigcup_{\alpha \in J} (X \setminus G_\alpha)$  is  $a$ - $m$ -open. Since  $X \setminus \bigcap_{\alpha \in J} G_\alpha = \bigcup_{\alpha \in J} (X \setminus G_\alpha)$ ,  $\bigcap_{\alpha \in J} G_\alpha$  is  $a$ - $m$ -closed.

**Remark 3.17** In a minimal structure space, by Corollary 3.16,  $aC_m(A)$  is  $a$ - $m$ -closed.

**Theorem 3.18** Let  $(X,m)$  be a minimal structure space and  $A \subseteq X$ . The following properties hold ;

- (1)  $aC_m(aC_m(A)) = aC_m(A)$ ,
- (2)  $aI_m(aI_m(A)) = aI_m(A)$ .

**Proof** (1) Clearly  $aC_m(A) \subseteq aC_m(aC_m(A))$ . Since  $aC_m(A)$  is  $a$ - $m$ -closed,  $aC_m(aC_m(A)) \subseteq aC_m(A)$ . Therefore  $aC_m(aC_m(A)) = aC_m(A)$ .

(2) Clearly  $aI_m(aI_m(A)) = aI_m(A)$ . Since  $aI_m(A)$  is  $a$ - $m$ -open,  $aI_m(A) \subseteq aI_m(aI_m(A))$ . Therefore  $aI_m(aI_m(A)) = aI_m(A)$ .

Let  $(X,m, \mathcal{I})$  be a minimal structure space with an ideal. For each  $x \in X$ , let  $\mathcal{M}^a(x) = \{U : x \in U, U \in \mathcal{M}^a\}$  be the family of all  $a$ - $m$ -open sets that contain  $x$ .

**Definition 3.19** Let  $(X,m, \mathcal{I})$  be a minimal structure space with an ideal and  $A \subseteq X$ . Then  $A_m^a(\mathcal{I}, m) = \{x \in X : U \cap A \notin \mathcal{I}, \text{ for every } U \in \mathcal{M}^a(x)\}$  is called  $a$ - $m$ -local function of  $A$  with respect to  $\mathcal{I}$  and  $m$ . We denote simply  $A_m^a$  for  $A_m^a(\mathcal{I}, m)$ .

**Remark 3.20** The minimal ideal is  $\{\emptyset\}$  and the maximal ideal is  $P(x)$  in any minimal structure space with an ideal  $(X,m, \mathcal{I})$ . It can be deduced that  $A_m^a(\{\emptyset\}, m) = aC_m(A)$  and  $A_m^a(P(x), m) = \emptyset$  for every  $A \subseteq X$ .

**Remark 3.21** In general,  $A \not\subseteq A_m^a$  and  $A_m^a \not\subseteq A$ . The next example shows that  $A \not\subseteq A_m^a$ .

**Example 3.22** Let  $X = \{a,b,c,d\}$  with a minimal structure  $m = \{\emptyset, \{a,b\}, \{b,c\}, \{c,d\}, \{a,d\}, X\}$  and  $\mathcal{I} = \{\emptyset, \{a\}, \{b\}, \{a,b\}\}$ ,  $A = \{a,b\}$ . Then  $\mathcal{M}^a = \{\emptyset, \{a,b\}, \{a,d\}, \{b,c\}, \{c,d\}, \{a,b,c\}, \{a,b,d\}, \{a,c,d\}, \{b,c,d\}, X\}$  and  $A_m^a = \emptyset$ .

**Theorem 3.23** Let  $(X,m, \mathcal{I})$  be a minimal structure space with an ideal and  $A, B \subseteq X$ . The following properties hold ;

- (1)  $(\emptyset)_m^a = \emptyset$ .
- (2) If  $A \subseteq B$ , then  $A_m^a \subseteq B_m^a$ .
- (3)  $(A_m^a)_m^a \subseteq A_m^a$ .
- (4)  $A_m^a \cup B_m^a \subseteq (A \cup B)_m^a$ .
- (5)  $(A \cap B)_m^a \subseteq A_m^a \cap B_m^a$ .
- (6)  $(A \setminus B)_m^a \setminus (B)_m^a \subseteq A_m^a \setminus B_m^a$ .

Proof (1) Assume  $(\emptyset)_m^a \neq \emptyset$ . Then there exists  $x \in (\emptyset)_m^a$ . Since  $X \in \mathcal{M}^a(X)$ ,  $X \cap \emptyset \notin \mathcal{I}$ . It contradicts with  $X \cap \emptyset = \emptyset \in \mathcal{I}$ . Therefore  $(\emptyset)_m^a = \emptyset$ .

(2) Assume that  $A \subseteq B$ . We will show that  $A_m^a \subseteq B_m^a$  by contrapositive. Suppose that  $x \notin B_m^a$ . Then there exists  $U \in \mathcal{M}^a(X)$  such that  $U \cap B \in \mathcal{I}$ . From  $A \subseteq B$  and the property of  $\mathcal{I}$ ,  $U \cap A \in \mathcal{I}$ . Therefore  $x \notin A_m^a$ .

(3) Assume that  $x \in (A_m^a)_m^a$ , and  $U \in \mathcal{M}^a(X)$ . Then  $A_m^a \cap U \notin \mathcal{I}$  and so  $A_m^a \cap U \neq \emptyset$ . Thus there exists  $y \in A_m^a \cap U$ , and so  $y \in U \in \mathcal{M}^a(y)$ . This implies that  $A \cap U \notin \mathcal{I}$ . Therefore  $x \in A_m^a$ .

(4) Since  $A \subseteq A \cup B$  and  $B \subseteq A \cup B$ , by (2)  $A_m^a \subseteq (A \cup B)_m^a$  and  $B_m^a \subseteq (A \cup B)_m^a$ . So  $A_m^a \cup B_m^a \subseteq (A \cup B)_m^a$ .

(5) Since  $A \cap B \subseteq A$  and  $A \cap B \subseteq B$ , by (2)  $(A \cap B)_m^a \subseteq A_m^a$  and  $(A \cap B)_m^a \subseteq B_m^a$ . So  $(A \cap B)_m^a \subseteq A_m^a \cap B_m^a$ .

(6) Since  $A \setminus B \subseteq A$ , by (2)  $(A \setminus B)_m^a \subseteq A_m^a$ . So  $(A \setminus B)_m^a \setminus B_m^a \subseteq A_m^a \setminus B_m^a$ .

Theorem 3.24 Let  $(X, m)$  be a minimal structure space and  $\mathcal{I}, \mathcal{J}$  are ideals on  $X$  where  $\mathcal{I} \subseteq \mathcal{J}$ . Then  $A_m^a(\mathcal{I}, m) \subseteq A_m^a(\mathcal{J}, m)$  for all  $A \subseteq X$ .

Proof Let  $A \subseteq X$ . Assume that  $x \in A_m^a(\mathcal{I}, m)$ . Then  $U \cap A \notin \mathcal{I}$  for every  $U \in \mathcal{M}^a(x)$ . Since  $\mathcal{I} \subseteq \mathcal{J}$ ,  $U \cap A \notin \mathcal{J}$  for every  $U \in \mathcal{M}^a(x)$ . Thus  $x \in A_m^a(\mathcal{J}, m)$ . Hence  $A_m^a(\mathcal{I}, m) \subseteq A_m^a(\mathcal{J}, m)$ .

Theorem 3.25 Let  $(X, m, \mathcal{I})$  be a minimal structure space with an ideal and  $A \subseteq X$ . The following properties hold ;

- (1)  $A_m^a \subseteq aC_m(A)$ ,
- (2)  $A_m^a = aC_m(A)$ , (i.e.,  $A_m^a$  is an  $a$ - $m$ -closed subset).

Proof (1) Assume that  $x \notin aC_m(A)$ . Then there exists an  $a$ - $m$ -closed set  $F$  such that  $A \subseteq F$  and  $x \notin F$ . Thus  $x \in X \setminus F$ , and so  $X \setminus F \in \mathcal{M}^a(x)$ . Hence  $(X \setminus F) \cap A = \emptyset \in \mathcal{I}$ , and so  $x \notin A_m^a$ . This implies that  $A_m^a \subseteq aC_m(A)$ .

(2) It is clear that  $A_m^a \subseteq aC_m(A_m^a)$ . Next, we will prove that  $aC_m(A_m^a) \subseteq A_m^a$ . Let  $x \in aC_m(A_m^a)$  and  $U \in \mathcal{M}^a(x)$ . Then  $A_m^a \cap U \neq \emptyset$ . Therefore there exists  $y \in A_m^a \cap U$ , so  $U \in \mathcal{M}^a(y)$ . Since  $y \in A_m^a$ ,  $A \cap U \notin \mathcal{I}$ , and so  $x \in A_m^a$ . Then  $A_m^a = aC_m(A_m^a)$ .

Theorem 3.26 Let  $(X, m, \mathcal{I})$  be a minimal structure space with an ideal and  $A \subseteq X$ . The following properties hold ;

- (1) If  $A \in \mathcal{I}$ , then  $A_m^a = \emptyset$ .
- (2) If  $U \in \mathcal{I}$ , then  $A_m^a = (A \cup U)_m^a$ .
- (3) If  $U \in \mathcal{I}$ , then  $A_m^a = (A \setminus U)_m^a$ .

Proof (1) Assume that  $A_m^a \neq \emptyset$ . Then there exists  $x \in A_m^a$ . Since  $X \in \mathcal{M}^a(x)$ ,  $A = X \cap A \in \mathcal{I}$ .

(2) Assume that  $U \in \mathcal{I}$ . Since  $A \subseteq A \cup U$  by Theorem 3.23(2), we get  $A_m^a \subseteq (A \cup U)_m^a$ . Next, we will prove that  $(A \cup U)_m^a \subseteq A_m^a$  by contrapositive. Suppose that  $x \notin A_m^a$ . Then there exists  $V \in \mathcal{M}^a(x)$  such that  $A \cap V \in \mathcal{I}$ . Since  $(A \cup U) \cap V = (A \cap V) \cup (U \cap V) \in \mathcal{I}$ ,  $(A \cup U) \cap V \in \mathcal{I}$ . Therefore  $x \notin (A \cup U)_m^a$ .

(3) Assume that  $U \in \mathcal{I}$ . Since  $A_m^a = (A \cap X)_m^a = (A \cap ((X \setminus U) \cup U))_m^a = ((A \setminus U) \cup (A \cap U))_m^a$  and  $A \cap U \subseteq U \in \mathcal{I}$ , by (2)  $A_m^a = (A \setminus U)_m^a$ .

Definition 3.27 Let  $(X, m, \mathcal{I})$  be a minimal structure with an ideal. An operator  $\mathfrak{R}_m^a : P(X) \rightarrow P(X)$  is defined as follows ; for every  $A \in P(X)$ ,  $\mathfrak{R}_m^a(A) = \{x \in X : \text{there exists } U \in \mathcal{M}^a(x) \text{ such that } U \setminus A \in \mathcal{I}\}$ .

Theorem 3.28 Let  $(X, m, \mathcal{I})$  be a minimal structure space with an ideal and  $A \in P(X)$ . Then  $\mathfrak{R}_m^a(A) = X \setminus (X \setminus A)_m^a$ .

Proof Let  $x \in \mathfrak{R}_m^a(A)$ . Then there exists an  $a$ - $m$ -open set  $U$  containing  $x$  such that  $U \setminus A \in \mathcal{I}$ . Thus  $U \cap (X \setminus A) \in \mathcal{I}$ . So  $x \notin (X \setminus A)_m^a$  and hence  $x \in X \setminus (X \setminus A)_m^a$ . Therefore  $\mathfrak{R}_m^a(A) \subseteq X \setminus (X \setminus A)_m^a$ .

For the reverse inclusion, let  $x \in X \setminus (X \setminus A)_m^a$ . Then  $x \notin (X \setminus A)_m^a$ . Thus there exists an  $a$ - $m$ -open set  $U$  containing  $x$  such that  $U \cap (X \setminus A) \in \mathcal{I}$ . This implies that  $U \setminus A \in \mathcal{I}$ . Hence  $x \in \mathfrak{R}_m^a(A)$ . So  $X \setminus (X \setminus A)_m^a \subseteq \mathfrak{R}_m^a(A)$ . Therefore  $\mathfrak{R}_m^a(A) = X \setminus (X \setminus A)_m^a$ .

Example 3.29 Let  $X = \{a, b, c, d\}$  with a minimal structure  $m = \{\emptyset, \{a, b\}, \{b, c\}, \{c, d\}, \{a, d\}, X\}$  and  $\mathcal{I} = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$ ,  $A = \{a, b\}$ . Then  $\mathfrak{R}_m^a(A) = \{\emptyset, \{a, b\}, \{a, d\}, \{b, c\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}, X\}$  and  $\mathfrak{R}_m^a(A) = \{a, b\}$ .

Theorem 3.30 Let  $(X, m, \mathcal{I})$  be a minimal structure space with an ideal, and  $A \subseteq X$ . Then  $\mathfrak{R}_m^a(A)$  is  $a$ - $m$ -open.



Proof We know that  $\mathfrak{R}_m^a(A) = X \setminus (X \setminus A)_m^a$  and  $(X \setminus A)_m^a$  is *a-m-closed*. Therefore  $\mathfrak{R}_m^a(A)$  is *a-m-open*.

Theorem 3.31 Let  $(X, m, \mathcal{S})$  be a minimal structure space with an ideal and  $A, B \subseteq X$ . Then the following properties hold ;

- (1) If  $A \subseteq B$ , then  $\mathfrak{R}_m^a(A) \subseteq \mathfrak{R}_m^a(B)$ .
- (2) If  $A \subseteq B$ , then  $\mathfrak{R}_m^a(A \cap B) \subseteq \mathfrak{R}_m^a(A) \cap \mathfrak{R}_m^a(B)$ .
- (3) If  $A \subseteq B$ , then  $\mathfrak{R}_m^a(A) \cup \mathfrak{R}_m^a(B) \subseteq \mathfrak{R}_m^a(A \cup B)$ .
- (4) If  $A \in \mathcal{M}^a$ , then  $A \subseteq \mathfrak{R}_m^a(A)$ .
- (5) If  $A \subseteq B$ , then  $\mathfrak{R}_m^a(A) \subseteq \mathfrak{R}_m^a(\mathfrak{R}_m^a(A))$ .

Proof (1) Assume that  $A \subseteq B$ . Then  $X \setminus B \subseteq X \setminus A$ . By Theorem 3.23(2),  $(X \setminus B)_m^a \subseteq (X \setminus A)_m^a$  and hence  $X \setminus (X \setminus A)_m^a \subseteq X \setminus (X \setminus B)_m^a$ . Therefore  $\mathfrak{R}_m^a(A) \subseteq \mathfrak{R}_m^a(B)$ .

(2) Since  $A \cap B \subseteq A$  and  $A \cap B \subseteq B$ ,  $\mathfrak{R}_m^a(A \cap B) \subseteq \mathfrak{R}_m^a(A)$  and  $\mathfrak{R}_m^a(A \cap B) \subseteq \mathfrak{R}_m^a(B)$ . Therefore  $\mathfrak{R}_m^a(A \cap B) \subseteq \mathfrak{R}_m^a(A) \cap \mathfrak{R}_m^a(B)$ .

(3) Since  $A \subseteq A \cup B$  and  $B \subseteq A \cup B$ ,  $\mathfrak{R}_m^a(A) \subseteq \mathfrak{R}_m^a(A \cup B)$  and  $\mathfrak{R}_m^a(B) \subseteq \mathfrak{R}_m^a(A \cup B)$ . Therefore  $\mathfrak{R}_m^a(A) \cup \mathfrak{R}_m^a(B) \subseteq \mathfrak{R}_m^a(A \cup B)$ .

(4) Assume that  $A \in \mathcal{M}^a$ . Then  $X \setminus A$  is *a-m-closed*. By Theorem 3.25(1), we get that  $(X \setminus A)_m^a \subseteq aC_m(X \setminus A) = X \setminus A$ . Therefore  $A = X \setminus (X \setminus A) \subseteq X \setminus (X \setminus A)_m^a = \mathfrak{R}_m^a(A)$ .

(5) By Theorem 3.30, we get that  $\mathfrak{R}_m^a(A)$  is *a-m-open*. By (4), we get that  $\mathfrak{R}_m^a(A) \subseteq \mathfrak{R}_m^a(\mathfrak{R}_m^a(A))$ .

Theorem 3.32 Let  $(X, m, \mathcal{S})$  be a minimal structure space with an ideal and  $A, B, U \subseteq X$ . Then the following properties hold ;

- (1) If  $U \in \mathcal{S}$ , then  $\mathfrak{R}_m^a(A \setminus U) = \mathfrak{R}_m^a(A)$ .
- (2) If  $U \in \mathcal{S}$ , then  $\mathfrak{R}_m^a(A \cup U) = \mathfrak{R}_m^a(A)$ .
- (3) If  $(A \setminus B) \cup (B \setminus A) \in \mathcal{S}$ , then  $\mathfrak{R}_m^a(A) = \mathfrak{R}_m^a(B)$ .
- (4) If  $A \in \mathcal{S}$ , then  $\mathfrak{R}_m^a(A) = X \setminus X_m^a$ .

Proof (1) Assume that  $A \subseteq X, U \in \mathcal{S}$ . By Theorem 3.26(2) and 3.28, we have  $\mathfrak{R}_m^a(A \setminus U) = X \setminus (X \setminus (A \setminus U))_m^a = X \setminus ((X \setminus A) \cup U)_m^a = X \setminus (X \setminus A)_m^a$ . Therefore  $\mathfrak{R}_m^a(A \setminus U) = \mathfrak{R}_m^a(A)$ .

(2) Assume that  $U \in \mathcal{S}$ . By Theorem 3.26(3), we have  $\mathfrak{R}_m^a(A \cup U) = X \setminus (X \setminus (A \cup U))_m^a = X \setminus ((X \setminus A) \setminus U)_m^a = X \setminus (X \setminus A)_m^a = \mathfrak{R}_m^a(A)$ .

(3) Assume that  $(A \setminus B) \cup (B \setminus A) \in \mathcal{S}$ .

Thus

$$\begin{aligned} \mathfrak{R}_m^a(A) &= \mathfrak{R}_m^a(A \setminus (A \setminus B)) \\ &= \mathfrak{R}_m^a((A \setminus (A \setminus B)) \cup (B \setminus A)) \\ &= \mathfrak{R}_m^a(B). \end{aligned}$$

(4) Assume that  $A \in \mathcal{S}$ . By Theorem 3.26(3), we get that  $\mathfrak{R}_m^a(A) = X \setminus (X \setminus A)_m^a = X \setminus X_m^a$ .

Theorem 3.33 Let  $(X, m, \mathcal{S})$  be a minimal structure space with an ideal and  $A \subseteq X$ . Then  $\mathfrak{R}_m^a(A) = \mathfrak{R}_m^a(\mathfrak{R}_m^a(A))$  if and only if  $(X \setminus A)_m^a = ((X \setminus A)_m^a)_m^a$ .

Proof It follows from the facts that,

$$\begin{aligned} \text{I) } \mathfrak{R}_m^a(A) &= X \setminus (X \setminus A)_m^a \text{ and } \mathfrak{R}_a \\ \text{II) } \mathfrak{R}_m^a(\mathfrak{R}_m^a(A)) &= X \setminus [X \setminus (X \setminus (X \setminus A)_m^a)]_m^a \\ &= X \setminus ((X \setminus A)_m^a)_m^a. \end{aligned}$$

Therefore  $\mathfrak{R}_m^a(A) = \mathfrak{R}_m^a(\mathfrak{R}_m^a(A))$  if and only if  $(X \setminus A)_m^a = ((X \setminus A)_m^a)_m^a$ .

### Discussion and Conclusion

The aim of this article is to introduce the results of properties of some sets in a minimal structure space with an ideal. In addition, we study some properties of  $\delta$ -*m-open* sets, *a-m-open* sets in a minimal structure space with an ideal. Moreover, we define an  $\delta$ -*m-local* function and an  $R_m^a$ -operator in a minimal structure space with an ideal. Some properties of them are obtained.

### Acknowledgements

The authors would like to thank the referees for helpful comments suggestions on the manuscript. It has received help from seniors, juniors, and friends in mathematics as well.

### References

Vaidyanathaswamy, R. (1945). The localization theory in set-topology. Proc. Indian Acad. Sci. 20, 51-61.  
 Al-Omeri, W, Noorani, M. & Al-Omari, A. (2014). a-local function and it's properties in ideal topological space. Fasc. Math, 53, 1-15.  
 Al-Omeri, W., Noorani, M. & Al-Omari, A. (2016). The operator in ideal topological space. Creat. Math, 25, 1-10.

- Noiri, T. & Popa, V. (2009). A generalization of some forms of g-irresolute functions. *European j. of Pure and Appl. Math*, 2(4): 473-493.
- Maki, H., Nagoor & Gani, K.C. (1999). On generalized semi-open and preopen sets. *Pure Appl. Math*, 49, 17-29.
- Rosas, E., Rajesh, N. & Carpintero, C. (2009). Some new type of open and closed sets in minimal structure-II. *International Mathematical Form*, 44, 2185-2198.
- Ozbakir, O.B. & Yildirim, E.D. (2009). On some closed sets in ideal minimal spaces. *Acta Math. Hungar*, 125(3), 227-235.
- Maki, H, Umehara, T. & Noiri, T. (1996). Every topological space in pre  $T_{1/2}$ . *Mem. Fac. Sci. Kochi Univ. Ser. A Math*. 17, 33-42.