

# ลำดับใหม่ที่สอดคล้องกับลำดับ $k$ -ฟีโบนัชชี

## Some novel sequences related to $k$ -Fibonacci sequences

พงษ์พันธ์ มุขวะชชี<sup>1</sup>, เพ็ญศิริ สมพงษ์<sup>2</sup>, สุพรรณณี สมพงษ์<sup>3</sup> และ ไกรลิขิต ลาตปะละ<sup>4\*</sup>

Phongphan Mukwachi<sup>1</sup>, Pensiri Sompong<sup>2</sup>, Supunnee Sompong<sup>3</sup> and Krailikhit Latpala<sup>4\*</sup>

Received: 13 March 2023 ; Revised: 18 May 2023 ; Accepted: 27 June 2023

### บทคัดย่อ

งานวิจัยนี้เราได้นำเสนอสามลำดับรูปแบบใหม่ของ  $\gamma_n$ ,  $\alpha_n$  และ  $\beta_n$  ที่มีส่วนเกี่ยวข้องกันผ่านความสัมพันธ์เวียนเกิด และเราได้สังเกตถึงความสัมพันธ์ของลำดับทั้งสามนี้สามารถแสดงให้อยู่ในรูปของลำดับ  $k$ -ฟีโบนัชชี เพื่อพิสูจน์ความสัมพันธ์นี้เราได้นำหลักอุปนัยเชิงคณิตศาสตร์ มาใช้สำหรับแสดงความถูกต้องของทฤษฎี และแสดงผลลัพธ์ที่ได้จากการศึกษาในงานนี้

คำสำคัญ: ลำดับ  $k$ -ฟีโบนัชชี, ความสัมพันธ์เวียนเกิด, อุปนัยเชิงคณิตศาสตร์

### Abstract

In this research, we introduce three novel sequences of  $\gamma_n$ ,  $\alpha_n$  and  $\beta_n$ . These sequences are related to each other through the recurrence relation, and we have observed that their relationship can be expressed using  $k$ -Fibonacci sequences. To prove this relationship, we used mathematical induction. We have shown the validity of our theorem, and the results are presented in this study.

**Keywords:**  $k$ -Fibonacci sequences, recurrence relation, mathematical induction

<sup>1</sup> อาจารย์ ภาควิชาคณิตศาสตร์และสถิติ คณะวิทยาศาสตร์และเทคโนโลยี มหาวิทยาลัยราชภัฏสกลนคร สกลนคร 47000

<sup>2</sup> ผู้ช่วยศาสตราจารย์ ภาควิชาวิทยาศาสตร์ทั่วไป คณะวิทยาศาสตร์และเทคโนโลยี มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตเฉลิมพระเกียรติ สกลนคร 47000

<sup>3</sup> ผู้ช่วยศาสตราจารย์ ภาควิชาคณิตศาสตร์และสถิติ คณะวิทยาศาสตร์และเทคโนโลยี มหาวิทยาลัยราชภัฏสกลนคร สกลนคร 47000

<sup>4</sup> อาจารย์ ภาควิชาคณิตศาสตร์ คณะครุศาสตร์ มหาวิทยาลัยราชภัฏสกลนคร สกลนคร 47000

<sup>1</sup> Lecturer, Department of Mathematics and Statistics, Faculty of Science and Technology, Sakon Nakhon Rajabhat University, Sakon Nakhon, 47000 Thailand

<sup>2</sup> Assistant Professor, Department of General Science, Faculty of Science and Engineering, Kasetsart University Chalermphrakiat Sakon Nakhon Province Campus, Sakon Nakhon 47000 Thailand

<sup>3</sup> Assistant Professor, Department of Mathematics and Statistics, Faculty of Science and Technology, Sakon Nakhon Rajabhat University, Sakon Nakhon, 47000 Thailand

<sup>4</sup> Lecturer, Department of Mathematics, Faculty of Education, Sakon Nakhon Rajabhat University, Sakon Nakhon, 47000 Thailand

\* Corresponding author: Krailikhit@snru.ac.th

**Introduction**

For any integer number  $k \geq 1$ , the  $n$  th  $k$ -Fibonacci sequence, denoted as  $\{F_{k,n}\}_{n=0}^\infty$ , is defined by (Falcon & Plaza, 2007) as a recursive sequence as follows:

$$F_{k,n+1} = kF_{k,n} + F_{k,n-1}$$

where  $F_{k,0} = 0$  and  $F_{k,1} = 1$ . The first 8 members of  $k$ -Fibonacci sequences are shown below:

$$0, 1, k, k^2 + 1, \dots, k^3 + 2k, k^4 + 3k^2 + 1, k^5 + 4k^3 + 3k, k^6 + 5k^4 + 6k^2 + 1.$$

(Atanassov, 2018) studied two new combined 3-Fibonacci sequences. Let  $a, b, c, d$  be arbitrary real numbers and  $\{F_n\}_{n=0}^\infty$  be the standard Fibonacci sequence. The first set of sequences has the form for  $n \geq 0$ ,

$$\alpha_{n+2} = \gamma_{n+1} + \beta_{n+1},$$

$$\beta_{n+2} = \gamma_{n+1} + \alpha_{n+1},$$

$$\gamma_{n+2} = \gamma_{n+1} + \gamma_n.$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \gamma_1 = d$ . From these sequences and for each natural number  $n \geq 1$  the result are the following,

$$\alpha_{2n+1} = b + F_{2n-1}a + (F_{2n}-1)d,$$

$$\alpha_{2n} = a + F_{2n}c + (F_{2n+1}-1)d,$$

$$\beta_{2n-1} = a + F_{2n-1}c + (F_{2n}+1)d,$$

$$\beta_{2n} = b + F_{2n}c + (F_{2n+1}-1)d,$$

$$\gamma_{n+2} = F_{n+1}c + F_{n+2}d.$$

The second set of sequences has the form for  $n \geq 0$ ,

$$\alpha_{n+1} = \alpha_{n+1} + \alpha_n,$$

$$\beta_{n+1} = \alpha_{n+1} + \gamma_n,$$

$$\gamma_{n+1} = \alpha_{n+1} + \beta_n.$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \alpha_1 = d$ . From these sequences and for each natural number  $n \geq 1$  the result are the following,

$$\alpha_n = F_{n-1}c + F_n d,$$

$$\beta_{2n-1} = (F_{2n}-1)a + b + (F_{2n+1}-1)d,$$

$$\beta_{2n} = (F_{2n+1}-1)a + c + (F_{2n+2}-1)d,$$

$$\gamma_{2n-1} = (F_{2n}-1)a + c + (F_{2n+1}-1)d,$$

$$\gamma_{2n} = (F_{2n+1}-1)a + b + (F_{2n+2}-1)d.$$

In the same year, he studied two additional new combined 3-Fibonacci sequences part 2. Let  $a, b, c$  be arbitrary real numbers and  $\{F_n\}_{n=0}^\infty$  be the standard Fibonacci sequence. The first set of sequences has the form for  $n \geq 0$ ,

$$\alpha_{n+1} = \beta_n + \gamma_n,$$

$$\beta_{n+1} = \alpha_n + \gamma_n,$$

$$\gamma_{n+1} = \frac{\alpha_{n+1} + \beta_{n+1}}{2} + \gamma_n,$$

where  $\alpha_0 = 2a, \beta_0 = 2b, \gamma_0 = c$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\alpha_n = (F_{2n-1}+(-1)^n)a + (F_{2n-1}-(-1)^n)b + F_{2n}c,$$

$$\beta_n = (F_{2n-1}-(-1)^n)a + (F_{2n-1}+(-1)^n)b + F_{2n}c,$$

$$\gamma_n = F_{2n}a + F_{2n}b + F_{2n+1}c.$$

The second set of sequences has the form for  $n \geq 0$ ,

$$\alpha_{n+1} = \alpha_n + \frac{\beta_n + \gamma_n}{2},$$

$$\beta_{n+1} = \alpha_{n+1} + \gamma_n,$$

$$\gamma_{n+1} = \alpha_{n+1} + \beta_n.$$

where  $\alpha_0 = a, \beta_0 = 2b, \gamma_0 = 2c$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\alpha_n = F_{2n-1}a + F_{2n}b + F_{2n}c,$$

$$\beta_n = F_{2n}a + (F_{2n+1} + (-1)^n)b + (F_{2n+1} - (-1)^n)c,$$

$$\gamma_n = F_{2n}a + (F_{2n+1} - (-1)^n)b + (F_{2n+1} + (-1)^n)c.$$

(Nubpetchploy & Pakapongpun, 2021) generated three combined sequences related to Jacobsthal sequences. Let  $a, b, c, d$  be arbitrary real numbers and  $J_n$  be the Jacobsthal sequences. The first set of sequences has the form for  $n \geq 0$ ,

$$\gamma_{n+2} = \gamma_{n+1} + 2\gamma_n,$$

$$\alpha_{n+1} = \gamma_{n+1} + 2\beta_n,$$

$$\beta_{n+1} = \gamma_{n+1} + 2\alpha_n.$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \gamma_1 = d$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\begin{aligned} \gamma_n &= 2J_{n-1}c + J_n d, \\ \alpha_n &= 2\alpha_{n-1} + (J_n + (-1)^n)c + j_n d + (-2)^n(a-b), \\ \beta_n &= 2\beta_{n-1} + (J_n + (-1)^n)c + j_n d - (-2)^n(a-b). \end{aligned}$$

The second set of sequences has the form for  $n \geq 0$ ,

$$\begin{aligned} \gamma_{n+2} &= \gamma_{n+1} + 2\gamma_n, \\ \alpha_{n+1} &= \gamma_n + 2\beta_n, \\ \beta_{n+1} &= \gamma_n + 2\alpha_n. \end{aligned}$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \gamma_1 = d$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\begin{aligned} \gamma_n &= 2J_{n-1}c + J_n d, \\ \alpha_n &= 2\alpha_{n-1} + (J_{n-1} + (-1)^{n-1})c + j_{n-1}d + (-2)^n(a-b), \\ \beta_n &= 2\beta_{n-1} + (J_{n-1} + (-1)^{n-1})c + j_{n-1}d - (-2)^n(a-b). \end{aligned}$$

The third set of sequences has the form for  $n \geq 0$ ,

$$\begin{aligned} \gamma_{n+1} &= \frac{\alpha_{n+1} + \beta_{n+1}}{2} + 2\gamma_n, \\ \alpha_{n+1} &= \gamma_n + 2\beta_n, \\ \beta_{n+1} &= \gamma_n + 2\alpha_n. \end{aligned}$$

where  $\alpha_0 = 2a, \beta_0 = 2b, \gamma_0 = c$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\begin{aligned} \gamma_{n-1} &= (J_{2n-1}-1)(a+b) + J_{2n-1}c, \\ \alpha_n &= (J_{n+1}^2 - J_n^2 + 1)(a+b) + (-1)^n J_n a + (-1)^{n+1} (2J_{n+1} \\ &+ J_n)b + J_{2n} c, \\ \beta_n &= (J_{n+1}^2 - J_n^2 + 1)(a+b) + (-1)^n J_n b + (-1)^{n+1} (2J_{n+1} \\ &+ J_n)a + J_{2n} c. \end{aligned}$$

(Atanassov, 2022) introduce on two new combined 3-Fibonacci sequences. Let  $a, b, c, d, e$  be arbitrary real numbers and  $\{F_n\}_{n=0}^\infty$  be the standard Fibonacci sequence. The first set of sequences has the form for  $n \geq 1$ ,

$$\begin{aligned} \alpha_{n+1} &= \alpha_n + \alpha_{n-1}, \\ \beta_{n+1} &= \beta_n + \beta_{n-1}, \\ \gamma_{n+1} &= \frac{\alpha_n + \beta_n}{2} + \gamma_n. \end{aligned}$$

where  $\alpha_0 = 2a, \beta_0 = 2b, \gamma_0 = c, \alpha_1 = 2d, \beta_1 = 2e$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\begin{aligned} \alpha_n &= 2F_{n-1}a + 2F_n d, \\ \beta_n &= 2F_{n-1}b + 2F_n e, \\ \gamma_n &= F_n a + F_n b + c + (F_{n+1}-1)d + (F_{n+1}-1)e. \end{aligned}$$

The second set of sequences has the form for  $n \geq 1$ ,

$$\begin{aligned} \alpha_{n+1} &= \alpha_n + \alpha_{n-1}, \\ \beta_{n+1} &= \beta_n + \beta_{n-1}, \\ \gamma_{n+1} &= \frac{\alpha_{n+1} + \beta_{n+1}}{2} + \gamma_n. \end{aligned}$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \alpha_1 = 2d, \beta_1 = 2e$ . From these sequences and for each natural number  $n \geq 1$  the result are the following.

$$\begin{aligned} \alpha_n &= 2F_{n-1}a + 2F_n d, \\ \beta_n &= 2F_{n-1}b + 2F_n e, \\ \gamma_n &= F_n a + F_n b + c + (F_{n+1}-1)d + (F_{n+1}-1)e. \end{aligned}$$

(Pakapongpun & Kongson, 2022) introduced three combined sequences related to  $k$ -Fibonacci sequences. Let  $a, b, c, d$  be arbitrary real numbers and  $\{F_{k,n}\}_{n=0}^\infty$  be the  $k$ -Fibonacci sequence. The first set of sequences has the form for  $n \geq 0$ ,

$$\begin{aligned} \gamma_{n+2} &= k\gamma_{n+1} + \gamma_n, \\ \alpha_{n+1} &= k\gamma_n + \beta_n, \\ \beta_{n+1} &= k\gamma_n + \alpha_n. \end{aligned}$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \gamma_1 = d$ . From these sequences the result are the following theorem 1.1.

Theorem 1.1. For any positive integer  $k$  and  $n$ ,

- (a)  $\gamma_n = F_{k,n}d + F_{k,n-1}c,$
- (b)  $\alpha_{2n} = (F_{k,2n} + F_{k,2n-1} - 1)d + (F_{k,2n-1} + F_{k,2n-2} + (F_{k,2} - 1)c + a,$
- (c)  $\beta_{2n} = (F_{k,2n} + F_{k,2n-1} - 1)d + (F_{k,2n-1} + F_{k,2n-2} + (F_{k,2} - 1)c + b,$
- (d)  $\alpha_{2n-1} = (F_{k,2n-1} + F_{k,2n-2} - 1)d + (F_{k,2n-2} + F_{k,2n-3} + (F_{k,2} - 1)c + b,$
- (e)  $\beta_{2n-1} = (F_{k,2n-1} + F_{k,2n-2} - 1)d + (F_{k,2n-2} + F_{k,2n-3} + (F_{k,2} - 1)c + a.$

The second set of sequences has the form for  $n \geq 0$ ,

$$\begin{aligned} \gamma_{n+2} &= k\gamma_{n+1} + \gamma_n, \\ \alpha_{n+1} &= k\gamma_{n+1} + \beta_n, \\ \beta_{n+1} &= k\gamma_{n+1} + \alpha_n. \end{aligned}$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c, \gamma_1 = d$ . From these sequences the result are the following theorem 1.2.

Theorem 1.2. For any positive integer  $k$  and  $n$ ,

- (a)  $\gamma_n = F_{k,n}d + F_{k,n-1}c,$
- (b)  $\alpha_{2n} = (F_{k,2n+1} + F_{k,2n} - 1)d + (F_{k,2n} + F_{k,2n-1} - 1)c + a,$
- (c)  $\beta_{2n} = (F_{k,2n+1} + F_{k,2n} - 1)d + (F_{k,2n} + F_{k,2n-1} - 1)c + b,$
- (d)  $\alpha_{2n-1} = (F_{k,2n} + F_{k,2n-1} - 1)d + (F_{k,2n-1} + F_{k,2n-2} - 1)c + b,$
- (e)  $\beta_{2n-1} = (F_{k,2n} + F_{k,2n-1} - 1)d + (F_{k,2n-2} + F_{k,2n-3} + (F_{k,2} - 1)c + a.$

The third set of sequences has the form for  $n \geq 0$ ,

$$\begin{aligned} \gamma_{n+1} &= k\gamma_n + \frac{\alpha_n + \beta_n}{2} \\ \alpha_{n+1} &= k\gamma_n + \beta_n, \\ \beta_{n+1} &= k\gamma_n + \alpha_n. \end{aligned}$$

where  $\alpha_0 = 2a, \beta_0 = 2b, \gamma_0 = c$ . From these sequences, the result are the following theorem 1.3.

Theorem 1.3. For any positive integer  $k$  and  $n$ ,

- (a)  $\gamma_{n+1} = \gamma_n (F_{k,2} + F_{k-1}) = \gamma_1 (F_{k,2} + F_{k-1})^n,$
- (b)  $\alpha_{2n} = \gamma_1 (F_{k,2} + F_{k-1})^{2n-1} + a - b,$
- (c)  $\alpha_{2n-1} = \gamma_1 (F_{k,2} + F_{k-1})^{2n-2} + b - a.$

In this paper, we introduce a new three set of combined sequences which are more general context related to  $k$ -Fibonacci sequences.

### Main Results

We applied those three sets of sequences from (Pakongpun & Kongson, 2022) work as follows. Let  $a, b, c, d$  and  $s$  be arbitrary real numbers with  $s \neq 0$ . The first set of sequences has the form for  $n \geq 0$ ,

$$\begin{aligned} \gamma_{n+2} &= k\gamma_{n+1} + \gamma_n, \\ \alpha_{n+1} &= ks\gamma_{n+1} + \beta_n, \\ \beta_{n+1} &= ks\gamma_{n+1} + \alpha_n. \end{aligned}$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c$  and  $\gamma_1 = d$ .

From these sequences, we generate the first few members of the sequences  $\{\gamma_n\}_{n=0}^\infty, \{\alpha_n\}_{n=0}^\infty$  and  $\{\beta_n\}_{n=0}^\infty$  with respect to  $n$  represented in Table 1, Table 2 and Table 3 respectively.

**Table 1** This table shows first 8 members of  $\{\gamma_n\}_{n=0}^\infty$  from the first set of sequences.

$n$	$\{\gamma_n\}_{n=0}^\infty$
0	$c$
1	$d$
2	$kd + c$
3	$k^2d + kc + d$
4	$k^3d + k^2c + c + 2kd + c$
5	$k^4d + k^3c + 3k^2d + 2kc + d$
6	$k^5d + k^4c + k^3d + 3k^2c + 3kd + c$
7	$k^6d + k^5c + 5k^4d + 4k^3c + 6k^2d + 3kc + d$

**Table 2** This table shows first 8 members of  $\{\alpha_n\}_{n=0}^\infty$  from the first set of sequences.

$n$	$\{\alpha_n\}_{n=0}^\infty$
0	$a$
1	$ksc + b$
2	$ks(c+d) + a$
3	$k^2sd + ks(2c+d) + b$
4	$k^3sd + k^2s(c+d) + ks(2c+2d) + a$
5	$k^4sd + k^3s(c+d) + k^2s(c+3d) + ks(3c+2d) + b$
6	$k^5sd + k^4s(c+d) + k^3s(c+4d) + k^2s(3c+3d) + ks(3c+3d) + a$
7	$k^6sd + k^5s(c+d) + k^4s(c+5d) + k^3s(4c+4d) + k^2s(3c+6d) + ks(4c+3d) + b$

**Table 3** This table shows first 8 members of  $\{\beta_n\}_{n=0}^\infty$  from the first set of sequences.

$n$	$\{\beta_n\}_{n=0}^\infty$
0	$b$
1	$ksc + a$
2	$ks(c+d) + b$
3	$k^2sd + ks(2c+d) + a$
4	$k^3sd + k^2s(c+d) + ks(2c+2d) + b$
5	$k^4sd + k^3s(c+d) + k^2s(c+3d) + ks(3c+2d) + a$
6	$k^5sd + k^4s(c+d) + k^3s(c+4d) + k^2s(3c+3d) + ks(3c+3d) + b$
7	$k^6sd + k^5s(c+d) + k^4s(c+5d) + k^3s(4c+4d) + k^2s(3c+6d) + ks(4c+3d) + a$

Theorem 2.1. For any positive integer  $k$  and  $n$ ,

- (a)  $\gamma_n = F_{k,n}d + F_{k,n-1}c,$
- (b)  $\alpha_{2n} = (F_{k,2n} + F_{k,2n-1} - 1)sd + (F_{k,2n-1} + F_{k,2n-2} + F_{k,2n-3} - 1)sc + a,$
- (c)  $\beta_{2n} = (F_{k,2n} + F_{k,2n-1} - 1)sd + (F_{k,2n-1} + F_{k,2n-2} + F_{k,2n-3} - 1)sc + b,$
- (d)  $\alpha_{2n-1} = (F_{k,2n-1} + F_{k,2n-2} - 1)sd + (F_{k,2n-2} + F_{k,2n-3} + F_{k,2n-4} - 1)sc + b,$  for  $n \geq 2,$
- (e)  $\beta_{2n-1} = (F_{k,2n-1} + F_{k,2n-2} - 1)sd + (F_{k,2n-2} + F_{k,2n-3} + F_{k,2n-4} - 1)sc + a,$  for  $n \geq 2.$

*Proof.* we will prove (a) by mathematical induction.

Let  $P(n)$  be a statement  $\gamma_n = F_{k,n}d + F_{k,n-1}c$  for  $n \geq 1$ , we will show that  $P(1)$  is true.

Since  $F_{k,1}d + F_{k,0}c = (1)d + (0)c = d = \gamma_1$ , then  $P(1)$  is true. Let  $m \geq 1$ , assume that  $P(1), P(2), \dots, P(m-1), P(m)$  are true that is,  $\gamma_n = F_{k,i}d + F_{k,i-1}c$ , where  $1 \leq i \leq m$ .

We will show that  $P(m+1)$  is true.

consider,

$$\begin{aligned} \gamma_{m+1} &= k\gamma_m + \gamma_{m-1} \\ &= k(F_{k,m}d + F_{k,m-1}c) + F_{k,m-1}d + F_{k,m-2}c \\ &= k(F_{k,m} + F_{k,m-1})d + (kF_{k,m-1} + F_{k,m-2})c \\ \gamma_{m+1} &= F_{k,m+1}d + F_{k,m}c. \end{aligned}$$

Then  $P(m+1)$  is true.

By mathematical induction, the statement  $P(n)$  is true for all  $n \geq 1$ .

Next, we will prove (b) by mathematical induction.

Let  $P(n)$  be a statement,

$$\begin{aligned} \alpha_{2n} &= (F_{n-2n} + F_{k,2n-1} - 1)sd \\ &\quad + (F_{k,2n-1} + F_{k,2n-2} + F_{k,2} - 1)sc + a, \text{ for } n \geq 1. \end{aligned}$$

We will show that  $P(1)$  is true.

Now consider,

$$\begin{aligned} &(F_{k,2(n)} + F_{k,2(1)-1} - 1)sd \\ &+ (F_{k,2(1)-1} + F_{k,2(1)-2} + F_{k,2} - 1)sc + a \\ &= (F_{k,2} + F_{k,1} - 1)sd + (F_{k,1} + F_{k,0} + F_{k,2} - 1)sc + a \\ &= (k+1-1)sd + (1+0+k-1)sc + a \\ &= ksd + ksc + a \\ &= ks(c+d) + a = \alpha_{2(1)}. \end{aligned}$$

Then  $P(1)$  is true.

Let  $m \geq 1$ , assume that  $P(m)$  is true that is,

$$\begin{aligned} \alpha_{2n} &= (F_{k-2m} + F_{k,2n-1} - 1)sd \\ &\quad + (F_{k,2m-1} + F_{k,2m-2} + F_{k,2} - 1)sc + a. \end{aligned}$$

We will show that  $P(m+1)$  is true.

Consider,

$$\begin{aligned} \alpha_{2m+2} &= ks\gamma_{2m+1} + \beta_{2m+1} \\ &= ks(F_{k,2m+1}d + F_{k,2m}c) + ks\gamma_{2m} + \alpha_{2m} \\ &= ks(F_{k,2m+1}d + F_{k,2m}c) + ks(F_{k,2m}d + F_{k,2m-1}c) \\ &\quad + (F_{k,2m} + F_{k,2m-1} - 1)sd + (F_{k,2m-1} + F_{k,2m-2} + \\ &\quad F_{k,2} - 1)sc + a \\ &= [(kF_{k,2m+1} + F_{k,2m})sd + (kF_{k,2m} + F_{k,2m-1}) \\ &\quad sd - sd] + [(kF_{k,2m} + F_{k,2m-1})sc + (kF_{k,2m-1} \\ &\quad + F_{k,2m-2})sc + F_{k,2}sc - sc] + a \\ &= (F_{k,2m+2} + F_{k,2m+1} - 1)sd + (F_{k,2m+1} + F_{k,2m} \\ &\quad + F_{k,2} - 1)sc + a \\ \alpha_{2(m+1)} &= (F_{k,2(m+1)} + F_{k,2(m+1)-1})sd \\ &\quad + (F_{k,2(m+1)-1} + F_{k,2(m+1)-2} + F_{k,2} - 1)sc + a. \end{aligned}$$

Then  $P(m+1)$  is true.

By mathematical induction the statement  $P(n)$  is true for all  $n > 1$ .

The proof of (c) is similar to (b).

To prove equation (d) for  $n \geq 2$ , using (a) and (c) we have,

$$\begin{aligned} \alpha_{2n-1} &= ks\gamma_{2n-2} + \beta_{2n-2} \\ &= ks(F_{k,2n-2}d + F_{k,2n-3}c) + (F_{k,2n-2} + F_{k,2n-3} - 1)sd \\ &\quad + (F_{k,2n-3} + F_{k,2n-4} + F_{k,2} - 1)sc + b \\ &= [(kF_{k,2n-2} + F_{k,2n-3})sd + (F_{k,2n-2}sd - sd)] + \\ &\quad [(kF_{k,2n-3} + F_{k,2n-4})sc + (F_{k,2n-3}sc + F_{k,2}sc - sc)] \\ &\quad + b \\ &= (F_{k,2n-1} + F_{k,2n-2} - 1)sd + (F_{k,2n-2} + F_{k,2n-3} \\ &\quad + F_{k,2} - 1)sc + b, \end{aligned}$$

then

$$\begin{aligned} \alpha_{2n-1} &= (F_{k,2n-1} + F_{k,2n-2} - 1)sd \\ &\quad + (F_{k,2n-2} + F_{k,2n-3} + F_{k,2} - 1)sc + b. \end{aligned}$$

is true.

**By (a), (b), and the proof is similar to (d), then we have (e).**

The proof is complete.

Next, we present the second sequences.

The second set of sequences has the form for

$n \geq 0$ ,

$$\begin{aligned} \gamma_{n+2} &= k\gamma_{n+1} + \gamma_n, \\ \alpha_{n+1} &= k\gamma_{n+1} + \beta_n, \\ \beta_{n+1} &= k\gamma_{n+1} + \alpha_n. \end{aligned}$$

where  $\alpha_0 = a, \beta_0 = b, \gamma_0 = c$  and  $\gamma_1 = d$ .

From these sequences, we generate the first 7 members of the sequences  $\{\alpha_n\}_{n=0}^\infty$  and  $\{\beta_n\}_{n=0}^\infty$  with respect to  $n$  represented in Table 4, and Table 5 respectively.

**Table 4** This table shows first 7 members of  $\{\alpha_n\}_{n=0}^\infty$  from the second set of sequences.

$n$	$\{\alpha_n\}_{n=0}^\infty$
0	$a$
1	$ksd + a$
2	$k^2sd + ks(c+d) + a$
3	$k^3sd + k^2s(c+d) + ks(c+2d) + b$
4	$k^4sd + k^3s(c+d) + k^2s(c+3d) + ks(2c+2d) + a$
5	$k^5sd + k^4s(c+d) + k^3s(c+4d) + k^2s(3c+3d) + ks(2c+3d) + b$
6	$k^6sd + k^5s(c+d) + k^4s(c+5d) + k^3s(4c+4d) + k^2s(3c+6d) + ks(3c+3d) + a$

**Table 5** This table shows first 7 members of  $\{\beta_n\}_{n=0}^\infty$  from the second set of sequences.

$n$	$\{\beta_n\}_{n=0}^\infty$
0	$b$
1	$ksd + a$
2	$k^2sd + ks(c+d) + b$
3	$k^3sd + k^2s(c+d) + ks(c+2d) + a$
4	$k^4sd + k^3s(c+d) + k^2s(c+3d) + ks(3c+2d) + b$
5	$k^5sd + k^4s(c+d) + k^3s(c+4d) + k^2s(3c+3d) + ks(2c+3d) + a$
6	$k^6sd + k^5s(c+d) + k^4s(c+5d) + k^3s(4c+4d) + k^2s(3c+6d) + ks(3c+3d) + b$

Theorem 2.2. For any positive integer  $k$  and  $n$ ,

- (a)  $\gamma_n = F_{k,n}d + F_{k,n-1}c,$
- (b)  $\alpha_{2n} = (F_{k,2n+1} + F_{k,2n} - 1)sd + (F_{k,2n} + F_{k,2n-1} - 1)sc + a,$
- (c)  $\beta_{2n} = (F_{k,2n+1} + F_{k,2n} - 1)sd + (F_{k,2n} + F_{k,2n-1} - 1)sc + b,$
- (d)  $\alpha_{2n-1} = (F_{k,2n} + F_{k,2n-1} - 1)sd + (F_{k,2n-1} + F_{k,2n-2} - 1)sc + b,$
- (e)  $\beta_{2n-1} = (F_{k,2n} + F_{k,2n-1} - 1)sd + (F_{k,2n-1} + F_{k,2n-2} - 1)sc + a.$

*Proof.* The proofs are similar to theorem 2.1.

Finally, the last sequences in our work.

The third set of sequences has the form for

$n \geq 0,$

$$\gamma_{n+1} = k\gamma_n + \frac{\alpha_n + \beta_n}{2s}$$

$$\alpha_{n+1} = ks\gamma_n + \beta_n,$$

$$\beta_{n+1} = ks\gamma_n + \alpha_n.$$

where  $\alpha_0 = 2as, \beta_0 = 2sb$  and  $\gamma_0 = c.$

The first 7 members of the sequences  $\{\gamma_n\}_{n=0}^\infty,$   
 $\{\alpha_n\}_{n=0}^\infty$  and  $\{\beta_n\}_{n=0}^\infty$  are show in Table 6, Table 7, and Table 8 respectively.

**Table 6** This table shows first 7 members of  $\{\gamma_n\}_{n=0}^\infty$  from the third set of sequences.

$n$	$\{\gamma_n\}_{n=0}^\infty$
0	$c$
1	$kc + a + b$
2	$k^2c + k(a+b+c) + a + b$
3	$k^3c + k^2(a+b+2c) + k(2a+2b+c) + a + b$
4	$k^4c + k^3(a+b+3c) + k^2(3a+3b+3c) + k(3c+3b+c) + a + b$
5	$k^5c + k^4(a+b+4c) + k^3(4c+4b+6c) + k^2(6a+6b+4c) + k(4a+4b+c) + a + b$
6	$k^6c + k^5(a+b+5c) + k^4(5a+5b+10c) + k^3(10a+10b+10c) + k^2(10a+10b+5c) + k(5a+5b+c) + a + b$

**Table 7** This table shows first 7 members of  $\{\alpha_n\}_{n=0}^\infty$  from the third set of sequences.

$n$	$\{\alpha_n\}_{n=0}^\infty$
0	$2as$
1	$ksc + 2bs$
2	$k^2sc + ks(a+b+c) + 2as$
3	$k^3sc + k^2s(a+b+2c) + ks(2a+2b+c) + 2bs$
4	$k^4sc + k^3s(a+b+3c) + k^2s(3a+3b+3c) + ks(3a+3b+c) + 2as$
5	$k^5sc + k^4s(a+b+4c) + k^3s(4c+4b+6c) + k^2s(6a+6b+4c) + ks(4a+4b+c) + 2bs$
6	$k^6sc + k^5s(a+b+5c) + k^4s(5a+5b+10c) + k^3s(10a+10b+10c) + k^2s(10a+10b+5c) + ks(5a+5b+c) + 2as$

**Table 8** This table shows first 7 members of  $\{\beta_n\}_{n=0}^\infty$  from the third set of sequences.

$n$	$\{\beta_n\}_{n=0}^\infty$
0	$2bs$
1	$ksc + 2as$
2	$k^2sc + ks(a+b+c) + 2bs$
3	$k^3sc + k^2s(a+b+2c) + ks(2a+2b+c) + 2as$
4	$k^4sc + k^3s(a+b+3c) + k^2s(3a+3b+3c) + ks(3a+3b+c) + 2bs$
5	$k^5sc + k^4s(a+b+4c) + k^3s(4c+4b+6c) + k^2s(6a+6b+4c) + ks(4a+4b+c) + 2as$
6	$k^6sc + k^5s(a+b+5c) + k^4s(5a+5b+10c) + k^3s(10a+10b+10c) + k^2s(10a+10b+5c) + ks(5a+5b+c) + 2bs$

Theorem 2.3. For any positive integer  $k$  and  $n$ ,

- (a)  $\gamma_{n+1} = \gamma_n(F_{k,2} + F_{k,1}) = \gamma_1(F_{k,2} + F_{k,1})^n,$
- (b)  $\alpha_{2n} = \gamma_1s(F_{k,2} + F_{k,1})^{2n-1} + as - bs,$
- (c)  $\beta_{2n} = \gamma_1s(F_{k,2} + F_{k,1})^{2n-1} + bs - as,$
- (d)  $\alpha_{2n-1} = \gamma_1s(F_{k,2} + F_{k,1})^{2n-2} + bs - as,$
- (e)  $\beta_{2n-1} = \gamma_1s(F_{k,2} + F_{k,1})^{2n-2} + as - bs.$

*Proof.* To prove (a) we will show that  $\gamma_{n+1} = \gamma_n(F_{k,2} + F_{k,1})$  since,  $\gamma_{n+1} = k\gamma_n + \frac{\alpha_n + \beta_n}{2s}$  and we know that,

$$\frac{\alpha_n + \beta_n}{2s} = \frac{(k\gamma_{n-1} + \beta_{n-1}) + (k\gamma_{n-1} + \alpha_{n-1})}{2s}$$

$$= k\gamma_{n-1} + \frac{\alpha_{n-1} + \beta_{n-1}}{2s},$$

so, we have  $\gamma_{n+1} = k\gamma_n + k\gamma_{n-1} + \frac{\alpha_{n-1} + \beta_{n-1}}{2s}$

Since  $\gamma_n = k\gamma_{n-1} + \frac{\alpha_{n-1} + \beta_{n-1}}{2s}$

we get that,

$$\begin{aligned} \gamma_{n+1} &= k\gamma_n + \gamma_n \\ &= \gamma_n(k+1) \end{aligned}$$

$$\gamma_{n+1} = \gamma_n(F_{k,2} + F_{k,1}).$$

Next, we will show that  $\gamma_{n+1} = \gamma_n(F_{k,2} + F_{k,1})^n$ .

Since  $\gamma_n = \gamma_{n-1}(F_{k,2} + F_{k,1})$

we have that,

$$\begin{aligned} \gamma_2 &= \gamma_1(F_{k,2} + F_{k,1}), \\ \gamma_3 &= \gamma_2(F_{k,2} + F_{k,1}) = \gamma_1(F_{k,2} + F_{k,1})^2, \\ \gamma_4 &= \gamma_3(F_{k,2} + F_{k,1}) = \gamma_1(F_{k,2} + F_{k,1})^3, \\ &\vdots \\ \gamma_{n+1} &= \gamma_1(F_{k,2} + F_{k,1})^n. \end{aligned}$$

thus  $\gamma_{n+1} = \gamma_n(F_{k,2} + F_{k,1}) = \gamma_1(F_{k,2} + F_{k,1})^n$ .

We will prove (b) by mathematical induction.

Let  $P(n)$  be the statement

$$\alpha_{2n} = \gamma_1 s(F_{k,2} + F_{k,1})^{2n-1} + as - bs \text{ for } n \geq 1.$$

We will show that  $P(1)$  is true.

consider,

$$\begin{aligned} &\gamma_1 s(F_{k,2} + F_{k,1})^{2(1)-1} + as - bs \\ &= s(kc+a+b)(k+1) + as - bs \\ &= k^2sc + ksa + ksb + ksc + as - bs + as + bs \\ &= k^2sc + ks(a+b+c) + 2as + \alpha_{2(1)} \end{aligned}$$

Then  $P(1)$  is true.

Let  $n \geq 1$ , assume that  $P(m)$  is true.

That is,  $\alpha_{2m} = \gamma_1 s(F_{k,2} + F_{k,1})^{2m-1} + as - bs$ .

We will show that  $P(m+1)$  is true.

Consider,

$$\begin{aligned} \alpha_{2(m+1)} &= \alpha_{2m+2} \\ &= ks\gamma_{2m+1} + \beta_{2m+1} \\ &= ks\gamma_{2m+1} + \beta_{2m} + \alpha_{2m} \\ &= ks\gamma_1(F_{k,2} + F_{k,1})^{2m} + ks\gamma_1(F_{k,2} + F_{k,1})^{2m-1} \\ &\quad + \gamma_1 s(F_{k,2} + F_{k,1})^{2m-1} + as - bs \\ &= ks\gamma_1(k+1)^{2m} + ks\gamma_1(k+1)^{2m-1} \\ &\quad + \gamma_1 s(k+1)^{2m-1} + as - bs \\ &= ks\gamma_1(k+1)(k+1)^{2m-1} + ks\gamma_1(k+1)^{2m-1} \\ &\quad + \gamma_1 s(k+1)^{2m-1} + as - bs \\ &= \gamma_1 s(k+1)^{2m-1} + [k(k+1)+k+1] + as - bs \\ &= \gamma_1 s(k+1)^{2m+1} + as - bs \\ &= \gamma_1 s(F_{k,2} + F_{k,1})^{2(m+1)-1} + as - bs \end{aligned}$$

then  $P(m+1)$  is true.

By mathematical induction the statement  $P(n)$  is true for all  $n \geq 1$ .

The proof of (c) is similar to (b).

From (a) and (c) we have (d), and similarly from (a) and (b) we also have (e).

### Conclusion and Discussion

A new three combined sequences related to  $k$ -Fibonacci sequences from new types were introduced and explicit formulas for their members are given.

From our sequences,

the first set of sequences,

$$\begin{aligned} \gamma_{n+2} &= k\gamma_{n+1} + \gamma_n \\ \alpha_{n+1} &= ks\gamma_n + \beta_n \\ \beta_{n+1} &= ks\gamma_n + \alpha_n \end{aligned}$$

the second set of sequences,

$$\begin{aligned} \gamma_{n+2} &= k\gamma_{n+1} + \gamma_n \\ \alpha_{n+1} &= ks\gamma_{n+1} + \beta_n \\ \beta_{n+1} &= ks\gamma_{n+1} + \alpha_n \end{aligned}$$



the third set of sequences,

$$\begin{aligned}\gamma_{n+1} &= k\gamma_n + \frac{\alpha_n + \beta_n}{2s} \\ \alpha_{n+1} &= ks\gamma_n + \beta_n, \\ \beta_{n+1} &= ks\gamma_n + \alpha_n.\end{aligned}$$

If  $s = 1$ , then the results correspond to the 3 set of sequences and the theorem 1.1, 1.2, and 1.3 in (Pakapongpun & Kongson, 2022). Other new schemes, modifying the standard form of  $k$ -Fibonacci sequences and new combined sequences will be discussed in the future.

### Acknowledgements

We thank the referees for their valuable reports that have improved this work.

### References

Falcon, S., & Plaza, Á. (2007). On the fibonacci k-numbers. *Chaos, Solitons Fractals*, 32(5), 1615-1624.

Atanassov, K. T. (2018). On two new combined 3-Fibonacci sequences. *Notes on Number Theory and Discrete Mathematics*, 24(2), 90-93.

Atanassov, K. T. (2018). On two new combined 3-Fibonacci sequences. Part 2. *Notes on Number Theory and Discrete Mathematics*, 24(3), 111-114.

Nubpetchploy, S., Pakapongpun, A. (2021). Three combined sequences related to Jacobsthal sequences. *International Journal of Mathematics and Computer Science*, 16(1), 149-157.

Atanassov, K. T. (2022). On two new combined 3-Fibonacci sequences. Part 3. *Notes on Number Theory and Discrete Mathematics*, 28(1), 143-146.

Pakapongpun, A., & Kongson, J. 2022. "Three combined sequences related to k-Fibonacci sequences. *Computer Science*, 7(2), 551-559.